

Sustainable management of *Pinus radiata* plantations



Cover photos:

Left: High pruning of radiata pine, New Zealand (P. Wilks)

Centre: A combination of radiata pine plantations, other introduced trees, native areas and farming create attractive landscapes in New Zealand; the farming is on the better soils (D. Mead)

Right: Recreation in a mature radiata pine plantation near Nelson, New Zealand (D. Mead)

Sustainable management of *Pinus radiata* plantations

FAO
FORESTRY
PAPER

170

by
Donald J. Mead

FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS
Rome 2013

Please cite as: **Mead, D.J.** 2013. Sustainable management of *Pinus radiata* plantations. FAO Forestry Paper No. 170. Rome, FAO.

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO.

ISBN 978-92-5-107634-7 (print)
E-ISBN 978-92-5-107635-4 (PDF)

© FAO 2013

FAO encourages the use, reproduction and dissemination of material in this information product. Except where otherwise indicated, material may be copied, downloaded and printed for private study, research and teaching purposes, or for use in non-commercial products or services, provided that appropriate acknowledgement of FAO as the source and copyright holder is given and that FAO's endorsement of users' views, products or services is not implied in any way.

All requests for translation and adaptation rights, and for resale and other commercial use rights should be made via www.fao.org/contact-us/licencerequest or addressed to copyright@fao.org.

FAO information products are available on the FAO website (www.fao.org/publications) and can be purchased through publications-sales@fao.org

All photographs and figures in this book are courtesy of the author, except where otherwise indicated.

Contents

Sustainable management of <i>Pinus radiata</i> plantations	i
Foreword	xii
Acknowledgements	xiii
Dedication	xiv
Acronyms and abbreviations	xv
1 Overview	1
GENERAL APPROACH	1
HISTORICAL PERSPECTIVE	3
The four phases of radiata pine plantation development	6
DISTRIBUTION OF RADIATA PINE PLANTATIONS	10
New Zealand	10
Chile	11
Australia	12
Spain	12
RADIATA PINE IN THE GLOBAL CONTEXT	12
2 Site requirements	15
MATCHING SPECIES TO SITE	15
CLIMATIC LIMITATIONS	16
Natural habitat	16
Exotic plantation experience	16
Other abiotic factors	19
Fire	22
Latitude and altitude	23
Aspect	23
Potential impacts of climate change	24
EDAPHIC LIMITATIONS	24
Physical properties of soil	25
Nutrient stresses	25
Diagnosing nutrient deficiencies	27
Soil microbiological factors	29
OTHER SITE CONSIDERATIONS	29
Catchment hydrology	30
RADIATA PINE'S ECOLOGICAL NICHE	32
3 Social, economic and environmental considerations	35
SOCIO-ECONOMIC SETTING	35
Australia	35
Chile	36
New Zealand	37
South Africa	38
Spain	39
Synthesis	39
ECONOMICS OF RADIATA PINE PLANTATIONS	40
Typical discount rates and plantation forest profitability	41

SOCIAL AND ECOSYSTEM SERVICES	43
Employment	44
Biodiversity	45
Landscape	48
Recreation in radiata pine plantations	48
Carbon storage	49
Wilding spread	49
Working with communities	50
ENVIRONMENTAL STANDARDS	50
TRENDS	51
4 Pests and diseases	53
MAJOR INSECT PESTS	53
Sirex wood wasp	53
Bark beetles	55
Pine shoot moth	56
Aphids and adelgids	56
Other localized insect problems	58
MAJOR DISEASES	58
Dothistroma needle blight	58
<i>Sphaeropsis sapinea</i>	60
Pine pitch canker	61
Other localized diseases	62
ANIMAL AND OTHER PESTS	63
PROSPECTS	63
5 Growth characteristics, wood properties and end-use	67
RADIATA PINE GROWTH	67
Growth habit	67
Shoot development	68
Growth stages	70
Growth patterns	71
Seasonal growth	72
Longer-term patterns	72
Productivity rating systems	74
Stocking and stand density	74
Crown growth	76
Branch development	77
Inter-tree competition and mortality	77
WOOD PROPERTIES AND END-USE	78
Cambial activity and differentiation	78
Corewood properties	79
Heartwood compared with sapwood	80
Basic density	80
Tracheids	81
Grain orientation and spiral grain	82
Compression wood	83
Knots and their link to grading systems	83
Clearwood	85
Log size and sweep	85
Other defects	85
Pulpwood and reconstituted products	86

OVERVIEW OF RADIATA PINE END USE	87
6 Radiata pine tree-breeding	91
IMPROVEMENT OBJECTIVES	91
BIOLOGICAL BACKGROUND	93
THE QUANTITATIVE GENETIC APPROACH	94
IMPROVEMENT OPTIONS	97
Provenance selection	97
Mass selection	98
Advanced breeding strategies	99
Incorporation of desired traits	102
Other recent developments	103
DOMESTICATION PROGRESS	103
7 Producing planting stock	105
THE SYSTEMS APPROACH AND SETTING OBJECTIVES	105
Optimum planting stock	106
SEED HANDLING	107
BARE-ROOTED PLANTING STOCK PRODUCTION	108
Seedbed preparation	109
Seed-sowing	109
Conditioning	110
Soil and nutrient management	112
Mycorrhizae	114
Irrigation	115
Control of nursery weeds, diseases and pests	115
VEGETATIVE PROPAGATION TECHNIQUES	116
CONTAINER PLANTS	119
PACKAGING AND TRANSPORT OF PLANTING STOCK	121
SYNTHESIS AND TRENDS	122
8 Establishment and early tending	125
ESTABLISHMENT PLANNING	125
THE BIOLOGICAL LIMITS TO EARLY GROWTH	127
SITE PREPARATION PRINCIPLES	130
RADIATA PINE SITE PREPARATION METHODS	131
Hand tool methods	131
Mechanical techniques	131
Fire	136
Chemicals	138
Other weed-control methods	139
PLANTING	140
Survival and replacements	141
DIRECT SEEDING AND NATURAL REGENERATION	142
FERTILIZER AT ESTABLISHMENT	143
FIRST ROTATION FEATURES	144
LATER ROTATION FEATURES	145
SYNTHESIS AND TRENDS	146

9 Tending established radiata pine stands	147
CHOICE OF TENDING SCHEDULES	147
Schedule evaluation	148
PRINCIPLES OF STAND DENSITY CONTROL	150
Initial stocking	151
Thinning objectives	152
Effect of stand density on stand characteristics	155
Other biotic and abiotic factors	156
Final crop stocking	157
Rotation length	158
NON-PRUNING TENDING SCHEDULES	159
Thinning techniques	161
PRINCIPLES OF PRUNING	162
Pruning objectives	163
PRUNING SCHEDULES	167
Pruning techniques	169
INTERRELATIONSHIPS AND FLEXIBILITY	169
Modelling systems	169
Schedule flexibility	170
USING RADIATA PINE IN MIXED SPECIES STANDS	170
FERTILIZERS	171
SYNTHESIS AND TRENDS	174
10 Productivity changes and sustainability of radiata pine plantation forests	177
CONCEPTS	177
PRODUCTIVITY INCREASES	179
PRODUCTIVITY DECREASES	182
Invasive species	182
Soil fertility changes	182
MANAGING INVASIVE SPECIES	185
MANAGING NUTRIENT SUSTAINABILITY	186
SYNTHESIS AND TRENDS	189
11 Radiata pine on farms	191
ECOLOGICAL PROCESSES IN RADIATA PINE SILVOPASTORAL SYSTEMS	191
Light competition	191
Moisture competition	193
Nutrient competition	193
Soil–plant interactions	194
Animal–plant interactions	194
Tree age effects	195
WINDBREAKS	196
WIDELY SPACED TREES OVER PASTURE	199
WOODLOTS	201
FARM PLANNING	202
TRENDS	202

12 Conclusions	205
LESSONS FROM THE RADIATA PINE EXPERIENCE	205
Growth characteristics	205
Species niche	205
Radiata pine forests and societal values and needs	206
Radiata pine's wood uses	206
Tree-breeding and silviculture	207
Sustainability	208
Uncertainties	209
THE FUTURE OF RADIATA PINE FORESTS	209
Glossary	211
References	215

List of Boxes

1.1 Early recognition of radiata pine's commercial potential	8
2.1 Radiata pine plantations reduce landslides	30
2.2 Longer-term effects of radiata pine plantations on stream flow	32
3.1 The new concept of using declining discount rates in forestry evaluations	41
4.1 The impact of parasites and hyperparasites on the control of the pine shoot moth, <i>Rhyacionia buoliana</i>	57
4.2 Pine pitch canker: a growing threat?	61
5.1 The effects of radiata pine wood density and stiffness on structural timber grades	82
6.1 The changing breeding goals for radiata pine: an example from Australasia	93
6.2 Structure of the joint New Zealand plus New South Wales radiata pine breeding programme	100
8.1 Managing available moisture for optimum growth of radiata pine in South Australia	134
9.1 Effect of silviculture on a fertile farm site in New Zealand	150
10.1 Second rotation radiata pine decline in South Australia reversed	185
11.1 Catchment diagnosis and design planning in New Zealand hill country	203

List of Figures

1.1	The plantation cycle, with major operations related to the planting stock production, establishment, stand tending and clearfelling of the crop	2
1.2	The Mt Peel radiata pine in Canterbury, New Zealand, planted in 1859 as a three-year-old seedling	3
1.3	Natural radiata pine stand at Monterey, California	5
1.4	Phases in the development of radiata pine plantation forestry	6
2.1	The interaction of factors affecting the choice of tree species	15
2.3	Drought deaths in a radiata pine stand in the Blackwood region of Western Australia	17
2.2	Radiata pine relative growth rate in relation to rainfall and soil depth	17
2.4	Severe defoliation due to abnormal climatic events in New Zealand causing physiological drought	18
2.5	Windthrow in a radiata pine stand the Nelson region of New Zealand	20
2.6	Stand edge windthrow following logging, which could have been avoided by careful planning	21
2.7	Sun-scald on radiata pine, Western Australia	21
2.8	Fire damage to young stand of radiata pine	22
2.9	Boron deficiency can cause repeated dieback	27
3.1	The New Zealand falcon (<i>Falco novaeseelandiae</i>) in a radiata pine plantation forest in Canterbury, New Zealand	45
3.2	Copihue, the Chilean national flower, growing in a radiata pine plantation forest in the BioBio region	46
3.3	A native forest remnant in radiata pine plantation forest in New Zealand	47
3.4	Recreation in mature radiata pine plantation forest near Auckland, New Zealand	48
4.1	The effects of the pine shoot moth, <i>Rhyaciona buoliana</i>	56
4.2	The effects of <i>Dothistroma</i> in radiata pine is usually most noticeable in the lower crown	59
4.3	<i>Sphaeropsis sapinea</i> (syn. <i>Diplodia pinea</i>) often results in leader dieback	60
4.4	Stem fluting caused by infection of pruned branches by <i>Neonectria fuckeliana</i>	62
5.1	Radiata pine shoot showing annual polycyclic growth pattern	67
5.2	Epicormic shoots forming on a pruned radiata pine tree, where the stem needles (short shoots) have not been removed	69
5.3	Generalized (not to scale) relationship between stocking, age (years) and growth	75
5.4	The effects of pruning and thinning on latewood percentage in South Africa	81
5.5	The relationship of branch index to structural grades	84
5.6	The impact of log small-end diameter and sweep on sawing recovery	86
5.7	Large radiata pine logs arriving at the mill	87
5.8	Radiata pine bark is widely used in horticulture	88
6.1	The use of selected clonal varieties can simultaneously improve both acoustic velocity (i.e. wood stiffness) and growth rate	96
6.2	Provenance differences: the bark on Monterey trees is thicker and more fissured than that on trees from Guadalupe Island	97
6.3	The hierarchy of populations	99

6.4	Control-pollinated seed orchards at Amberley, New Zealand	102
7.1	The effects of spacing within the nursery bed and seedling diameters at lifting, on volume index (D^2H) at the end of the first growing season after planting	107
7.2	Bare-rooted radiata plants are grown in long raised beds	109
7.3	Cuttings of radiata pine from stool beds set directly into beds and protected by plastic covers	117
7.4	One-year-old cuttings with a good diameter stem and a vigorous root system with mycorrhizae	118
7.5	Somatic embryogenesis plants in a misted greenhouse in Chile before being on-grown	119
7.6	Typical flowchart of growing clones of radiata pine, starting from control-pollinated seed, through laboratory multiplication and storage to nursery	120
7.7	Container grown radiata pine employing air-pruning of roots	121
7.8	Bare-rooted stock lifted and placed horizontally into boxes	122
7.9	Trimming the roots to 10 cm length	123
8.1	Factors affecting the stand at the end of the establishment period	126
8.2	Growth responses to silvicultural treatments	129
8.3	Site preparation in South Australia using towed rollers. The resulting mulch assists with moisture conservation, conserves nutrients and makes planting easier.	135
8.4	Mounding and ripping provide an excellent cultivated planting site, concentrate nutrients and reduce weed competition.	135
8.5	Windrows made using a backhoe with a root-rake attachment. Note that there is some soil disturbance.	136
8.6	The heaping of slash, and then burning the heaps, clears the site but runs the risk of nutrient loss and creating uneven growth in the compartment.	137
8.7	Container-planted seedling, illustrating the regeneration of new roots four months after planting. Note that part of the foliage had been deliberately buried.	141
8.8	Trees replaced in gaps seldom keep up with those planted earlier. The large trees are three years old.	142
8.9	Hauler logging leaves a clean site but redistributes nutrients.	145
9.1	The rise of green crowns results in the development of bark-encased knots after branches die	156
9.2	Mechanized thinning in South Australia	159
9.3	Stands of radiata pine with too high a height-diameter ratio can have stability problems	162
9.4	Ladder high pruning of radiata pine in New Zealand	163
9.5	Christmas trees trimmed to create dense, conical crowns	167
9.6	Long crowns left after pruning in Chile avoid suppression of the pruned trees by non-pruned trees	168
9.7	Phosphate fertilizer response on a deficient site: a site with fertilizer (left) and without (right)	172
10.1	Tree growth is determined by the most limiting factor and some of these factors can be manipulated by forest managers	178
10.2	The effect of slash burning (top) compared with slash retention (bottom) on radiata pine at age four years on sands in Western Australia	181

10.3	Overview of site assessment, level of monitoring in relation to harvesting impacts and the need for nutrient amendments	187
11.1	Pasture production under radiata pine trees, as influenced by final crop stocking, stand arrangement and site	192
11.2	Contour soil bund to collect runoff in a low-rainfall area in Chile	194
11.3	Radiata pine in Chile, at age 5 years, planted in twin rows spaced 7m apart. The ripping rows follow the contour to encourage water infiltration	195
11.4	The effects of windbreak porosity on windflow at different distances from the windbreak for porous, single-row and multiple-row conifers and for a solid barrier	197
11.5	Representation of a two-row shelterbelt layout, with supplementary species to windward	198
11.6	A porous two-row radiata pine and deodar cedar shelterbelt that has been side-trimmed and in which every second radiata pine tree has been high-pruned	198
11.7	A radiata pine shelterbelt designed to catch snow	199
12.1	Radiata pine logs on their way to the wood-using industry	207
12.2	Radiata pine, other introduced trees, native areas and farming create an attractive landscape	209

List of Tables

1.1	Timeline for the domestication of radiata pine and the development of silviculture	7
1.2	Estimated radiata pine plantation areas and the average annual change in area over the last five years, together with major uses by country	10
1.3	Percentage of stands for radiata pine by tending schedules and age class for the whole of New Zealand and for the Central North Island Region (CNI) in 2011	11
1.4	Growth rates, typical rotation lengths and areas of selected species used in large-scale plantations in selected countries	13
2.1	General climate profiles for radiata pine and <i>Pinus taeda</i> developed for models used to select regions to grow these species	19
2.2	Nutrient deficiency symptoms in radiata pine, marginal foliage nutrient concentrations and the confidence in these levels	28
3.1	Characteristics of the main countries with radiata pine plantations	35
3.2	Economic comparison of forest plantations of various species in selected countries	42
3.3	Ecosystem services provided by radiata pine plantation forests and associated agroforestry systems	43
5.1	The influence of stocking on tree size, green crown height, height-to-diameter ratio and stiffness (MoE) for the outer wood: results from a 17 year-old radiata pine Nelder spacing trial in New Zealand	76
6.1	Estimated heritabilities (h^2) for a range of radiata pine traits	95
6.2	Selected provenance differences in radiata pine	98
6.3	Improvement ratings for growth and form for different seedlots, New Zealand	99
7.1	Plant types and their key characteristics available for planting <i>Pinus radiata</i>	105
7.2	Specifications for bare-root radiata pine seedlings and cuttings*	106
7.3	Mean nutrient content in open-grown seedlings, at lifting	112
7.4	Characteristics of cuttings compared with seedlings of <i>Pinus radiata</i>	116
8.1	Steps in the establishment planning process	127
8.2	Fine root density of radiata pine and weeds in young stands growing in a sandy soil, South Australia	129
8.3	Site preparation methods commonly used in radiata pine plantations	132
9.1	The current range of typical radiata pine thinning schedules in non-pruned stands	153
9.2	Current typical pruning schedules with radiata pine	154
9.3	Typical fertilizer types and rates used in radiata pine plantations	173
10.1	Factors increasing radiata pine productivity, classified by response type and translated to gain at the end of a rotation	180
10.2	Typical nutrient balance sheets for selected contrasting sites and harvesting practices, New Zealand	183

Foreword

Growing societal demand for forest products and services require an increase in efforts to establish new forests. Where appropriate, intensively managed forest plantations can supply local and global markets with wood while contributing significantly to rural and industrial development. One of the most successful forest plantation species is *Pinus radiata* – radiata pine.

FAO recognizes the need to back up any significant tree-planting with a deep understanding of all aspects of the species being planted. In 1960, FAO published a major review of radiata pine, written by C.W. Scott, which presented the state of knowledge at that time. After more than 50 years, FAO is pleased to present a comprehensive update of knowledge on radiata pine. While this new publication, by Dr Donald Mead, covers many aspects of the species – its discovery and domestication, site requirements, limitations, wood properties and end-uses, and its social and environmental roles – its focus is on the principles and practices of growing radiata pine sustainably. It also looks ahead to emerging challenges facing plantation forest management, such as the effects of climate change, new diseases and other threats, and how to manage plantation forests to meet changing product needs and societal demands. Thus, this book is relevant and helpful for growers of other tree species, and should also provide valuable insights to forestry students and a wide range of other people interested in forestry.

Globally, there are just over 4 million ha of radiata pine plantations, mainly in Australia, Chile, New Zealand and Spain, making this species the most widely planted introduced conifer. As a plantation species, radiata pine has been a spectacular success, becoming the basis of strong wood-using industries in those countries and producing about 60 million cubic metres of wood per year. Even so, radiata pine provides only slightly more than four percent of the output of all planted forests.

Dr Mead, a New Zealander, has been involved in forestry research and teaching about radiata pine for more than 50 years. He wanted FAO to publish this review of radiata pine management as a means to ensuring its wide dissemination. For FAO, this publication is a way of educating professionals and others on the importance of long-term sustainable plantation forest management based on a profound knowledge of the biology, silviculture and use of a species. I congratulate Dr Mead for this outstanding contribution to global knowledge on plantation forest management and recommend the work to the global forestry community.



Eduardo Rojas-Briales
Assistant Director-General
FAO Forestry Department

Acknowledgements

This book has endured a long gestation. My sincere thanks go to the many people who provided information over the years. In particular, I thank Peter Lavery and Gonzalo Paredes, who were instrumental in providing the impetus for this book in the early years. Gonzalo supported the writing of this book until his health prevented him from being actively involved in the last few months.

I also sincerely thank Alvaro Sotomayor, Rowland Burdon, Euan Mason, Lisa Langer, Mike Carson, Eric Appleton, Robin Trewin, Piers Maclaren, Jose Prado, Tat Smith and Philip West, all of whom reviewed parts of this book. Rowland Burdon actively cooperated by providing background information on radiata pine history and status; he has been involved with a book on the domestication of the species. Jim Carle provided insightful comments on the entire book based on his wide experience with FAO and as a forest manager. Walter Kollert is also thanked for his support after Jim Carle retired. Rosamund Arthur assisted by reading the manuscript from a non-forester's viewpoint. I thank the editorial staff at FAO – Giulia Muir, Alastair Sarre and Patricia Tendi – and designer Roberto Cenciarelli for their excellent work.

Financial support for this project was provided by the New Zealand Forestry Corporation, the University of Canterbury and Lincoln University, the latter two by way of study leave.

Dedication

This book is dedicated to Dr Gonzalo Paredes Veloso, who died at Valdivia, Chile, in October 2012. Gonzalo was a forest engineer who did his PhD in forest economics at Corvallis, Oregon, United States of America, from 1984 to 1986. He was Dean of the Faculty of Forestry and Natural Resources, Universidad Austral de Chile in Valdivia, for two periods of his career, and for a time he was the Vice Chancellor of Finance and Administrative Support. He was a major leader in the development of the Chilean radiata pine simulator and actively supported forestry research in Chile. Between 1997 and 2000 he was director of Instituto Forestal, the Chilean government forest research organization.

Gonzalo, an academic leader and a critical thinker, was a strong supporter of this book project. He was a dedicated family man and a good friend to many, including the author.

Acronyms and abbreviations

B	boron
BA	basal area
C	carbon
Ca	calcium
CAI	current annual increment
CO ²	carbon dioxide
CSIRO	Commonwealth Scientific and Industrial Research Organisation (Australia)
Cu	copper
D	diameter
dbh	diameter at breast height
DCF	discounted cash flow
DDC	diameter of the defect core in the centre of a pruned log
DOO	diameter over occlusion
DOS	maximum stem diameter over branch stubs after pruning
DRIS	Diagnosis and Recommendation Integrated System
FAO	Food and Agriculture Organization of the United Nations
Fe	iron
FRI	Forest Research Institute (New Zealand), now Scion
FSC	Forest Stewardship Council
G	genetic gain
H	tree height
H ²	broad-sense heritability
INFOR	Instituto Forestal (Chile)
IRR	internal rate of return
LAI	leaf area index
LEV	land expectation value
LiDAR	light detection and ranging
LSW	log sweep
MAI	mean annual increment since establishment
Mg	magnesium
Mn	manganese
MoE	modulus of elasticity
N	nitrogen
n	stocking
NPW	net present worth
P	phosphorus
PEFC	Programme for the Endorsement of Forest Certification
PMAI	periodic mean annual increment
RD	relative density
REDD+	reducing emissions from deforestation and forest degradation
S	sulphur
SI	site index
V	variance
Zn	zinc