



Food and Agriculture Organization
of the United Nations

**FAO SPECIFICATIONS AND
EVALUATIONS FOR AGRICULTURAL
PESTICIDES**

PROPOXUR

2-isopropoxyphenyl methylcarbamate

TABLE OF CONTENTS

	PAGE
DISCLAIMER	
INTRODUCTION	1
PART ONE	
SPECIFICATIONS FOR PROPOXUR	
PROPOXUR INFORMATION	3
PROPOXUR TECHNICAL MATERIAL (OCTOBER 2021)	4
PROPOXUR DUSTABLE POWDER (OCTOBER 2021)	5
PROPOXUR WETTABLE POWDER (OCTOBER 2021)	7
PROPOXUR EMULSIFIABLE CONCENTRATE (OCTOBER 2021)	9
PART TWO	
EVALUATIONS OF PROPOXUR	11
2021 FAO/WHO EVALUATION REPORT ON PROPOXUR	12
SUPPORTING INFORMATION	14
ANNEX 1: HAZARD SUMMARY PROVIDED BY THE PROPOSER	16
ANNEX 2: REFERENCES	18
2016 FAO/WHO EVALUATION REPORT ON PROPOXUR	19
SUPPORTING INFORMATION	23
ANNEX 1: HAZARD SUMMARY PROVIDED BY THE PROPOSER	25
ANNEX 2: REFERENCES	27
2003 EVALUATION REPORT FOR PROPOXUR	28

DISCLAIMER¹

FAO specifications are developed with the basic objective of promoting, as far as practicable, the manufacture, distribution and use of pesticides that meet basic quality requirements.

Compliance with the specifications does not constitute an endorsement or warranty of the fitness of a particular pesticide for a particular purpose, including its suitability for the control of any given pest, or its suitability for use in a particular area. Owing to the complexity of the problems involved, the suitability of pesticides for a particular purpose and the content of the labelling instructions must be decided at the national or provincial level.

Furthermore, pesticides which are manufactured to comply with these specifications are not exempted from any safety regulation or other legal or administrative provision applicable to their manufacture, sale, transportation, storage, handling, preparation and/or use.

FAO disclaims any and all liability for any injury, death, loss, damage or other prejudice of any kind that may arise as a result of, or in connection with, the manufacture, sale, transportation, storage, handling, preparation and/or use of pesticides which are found, or are claimed, to have been manufactured to comply with these specifications.

Additionally, FAO wishes to alert users to the fact that improper storage, handling, preparation and/or use of pesticides can result in either a lowering or complete loss of safety and/or efficacy.

FAO is not responsible, and does not accept any liability, for the testing of pesticides for compliance with the specifications, nor for any methods recommended and/or used for testing compliance. As a result, FAO does not in any way warrant or represent that any pesticide claimed to comply with a FAO specification actually does so.

¹ This disclaimer applies to all specifications published by FAO.

INTRODUCTION

FAO establishes and publishes specifications* for technical material and related formulations of agricultural pesticides, with the objective that these specifications may be used to provide an international point of reference against which products can be judged either for regulatory purposes or in commercial dealings.

From 1999 onward, the development of FAO specifications follows the **New Procedure**, described first in the 5th edition of the "Manual on the development and use of FAO specifications for plant protection products" and later in the 1st edition of "Manual for Development and Use of FAO and WHO Specifications for Pesticides" (2002) - currently available as 3rd revision of the 1st edition (2016) - , which is available only on the internet through the FAO and WHO web sites.

This **New Procedure** follows a formal and transparent evaluation process. It describes the minimum data package, the procedure and evaluation applied by FAO and the Experts of the FAO/WHO Joint Meeting on Pesticide Specifications (JMPS). [Note: prior to 2002, the Experts were of the FAO Panel of Experts on Pesticide Specifications, Registration Requirements, Application Standards and Prior Informed Consent, which now forms part of the JMPM, rather than the JMPS.]

FAO Specifications now only apply to products for which the technical materials have been evaluated. Consequently from the year 1999 onwards the publication of FAO specifications under the **New Procedure** has changed. Every specification consists now of two parts namely the specifications and the evaluation report(s):

Part One: The Specification of the technical material and the related formulations of the pesticide in accordance with chapters 4 to 9 of the "Manual on development and use of FAO and WHO specifications for pesticides".

Part Two: The Evaluation Report(s) of the pesticide, reflecting the evaluation of the data package carried out by FAO and the JMPS. The data are provided by the manufacturer(s) according to the requirements of chapter 3 of the "FAO/WHO Manual on Pesticide Specifications" and supported by other information sources. The Evaluation Report includes the name(s) of the manufacturer(s) whose technical material has been evaluated. Evaluation reports on specifications developed subsequently to the original set of specifications are added in a chronological order to this report.

FAO specifications developed under the **New Procedure** do not necessarily apply to nominally similar products of other manufacturer(s), nor to those where the active ingredient is produced by other routes of manufacture. FAO has the possibility to extend the scope of the specifications to similar products but only when the JMPS has been satisfied that the additional products are equivalent to that which formed the basis of the reference specification.

Specifications bear the date (month and year) of publication of the current version. Evaluations bear the date (year) of the Meeting at which the recommendations were made by the JMPS.

* NOTE: PUBLICATIONS ARE AVAILABLE ON THE INTERNET AT (<http://www.fao.org/pest-and-pesticide-management/guidelines-standards/faowho-joint-meeting-on-pesticide-specifications-jmps/pesticide-specifications/pesticide-specifications-list/en/>) OR IN HARDCOPY FROM THE PLANT PROTECTION INFORMATION OFFICER.

PART ONE

SPECIFICATIONS

PROPOXUR	PAGE
PROPOXUR INFORMATION	3
PROPOXUR TECHNICAL MATERIAL (OCTOBER 2021)	4
PROPOXUR DUSTABLE POWDER (OCTOBER 2021)	5
PROPOXUR WETTABLE POWDER (OCTOBER 2021)	7
PROPOXUR EMULSIFIABLE CONCENTRATE (OCTOBER 2021)	9

PROPOXUR

INFORMATION

Common name

Propoxur (E-ISO, F-ISO)

Synonyms

Bay 9010, Baygon, Bayer 39007, Blattanex, Bolfo, BO Q 5812315, OMS 33, PHC (JMAF), Pillargon, UN Carbamate, Tugon, Unden, Undene.

Chemical names

IUPAC: 2-isopropoxyphenyl methylcarbamate

CA: 2-(1-methylethoxy)phenyl methylcarbamate


CAS Registry number

114-26-1

CIPAC number

80

Structural formula



Solid propoxur can exist in two crystal forms (modifications I and II) but the technical material usually contains >95% of modification I.

Empirical formula

C₁₁H₁₅NO₃

Relative molecular mass

209.25

Identity tests

HPLC retention time, with detection at 280 nm (CIPAC Handbook D, p. 155, 1988); IR and mass spectra; melting point (87.5-90°C).

PROPOXUR TECHNICAL MATERIAL

FAO Specification 80 / TC (October 2021*)

This specification, which is PART ONE of this publication, is based on an evaluation of data submitted by the manufacturers whose names are listed in the evaluation reports (80/2003, 80/2016 & 80/2021). It should be applicable to TC of these manufacturers but it is not an endorsement of those products, nor a guarantee that they comply with the specification. The specification may not be appropriate for TC produced by other manufacturers. The evaluation reports (80/2003, 80/2016 & 80/2021), as PART TWO, form an integral part of this publication.

1 Description

The material shall consist of propoxur together with related manufacturing impurities, in the form of colourless to pale yellow crystals with a phenolic odour, free from visible extraneous matter and added modifying agents.

2 Active ingredient

2.1 Identity tests (80/TC/M2/2, CIPAC Handbook D, p. 155, 1988)

The active ingredient shall comply with an identity test and, where the identity remains in doubt, shall comply with at least one additional test.

2.2 Propoxur content (80/TC/M2/3, CIPAC Handbook D, p. 155, 1988)

The propoxur content shall be declared (not less than 980 g/kg) and, when determined, the average measured content shall not be lower than the declared minimum content.

3 Relevant impurities

3.1 Water (MT 30.6, CIPAC Handbook P, p. 222, 2021)

Maximum: 2.0 g/kg.

3.2 Material insoluble in acetone (MT 27, CIPAC Handbook F, p. 88, 1995)

Maximum: 1.0 g/kg

4 Physical properties

4.1 Acidity and alkalinity (MT 31.2.1, CIPAC Handbook F, p. 98, 1995)

Maximum acidity: 0.5 g/kg calculated as H₂SO₄.

Maximum alkalinity: 0.1 g/kg calculated as NaOH.

* Specifications may be revised and/or additional evaluations may be undertaken. Ensure the use of current versions by checking at: <https://www.fao.org/pest-and-pesticide-management/guidelines-standards/faowho-joint-meeting-on-pesticide-specifications-jmps/pesticide-specifications/pesticide-specifications-list/en/>

PROPOXUR DUSTABLE POWDER

FAO Specification 80 / DP (October 2021*)

This specification, which is PART ONE of this publication, is based on an evaluation of data submitted by the manufacturer whose name is listed in the evaluation report 80/2003. It should be applicable to relevant products of this manufacturer but it is not an endorsement of those products, nor a guarantee that they comply with the specification. The specification may not be appropriate for the products of other manufacturers. The evaluation report 80/2003, as PART TWO, forms an integral part of this publication.

1 Description

The material shall consist of an homogeneous mixture of technical propoxur, complying with the requirements of FAO specification 80/TC (October 2021), together with any other necessary carriers and any other necessary formulants. It shall be in the form of a fine, free-flowing, beige powder, free from visible extraneous matter and hard lumps.

2 Active ingredient

2.1 Identity tests (80/TC/M2/2, CIPAC Handbook D, p. 155, 1988)

The active ingredient shall comply with an identity test and, where the identity remains in doubt, shall comply with at least one additional test.

2.2 Propoxur content (80/DP/M2/3, CIPAC Handbook D, p. 155, 1988)

The propoxur content shall be declared (g/kg) and, when determined, the average content measured shall not differ from that declared by more than the following tolerances.

Declared content in g/kg	Tolerance
Up to 25	± 15 % of the declared content
above 25 up to 100	± 10 % of the declared content
Note in each range the upper limit is included.	

3 Relevant impurities

3.1 Water (MT 30.6, CIPAC Handbook P, p. 222, 2021)

Maximum: 15 g/kg.

* Specifications may be revised and/or additional evaluations may be undertaken. Ensure the use of current versions by checking at: <https://www.fao.org/pest-and-pesticide-management/guidelines-standards/faowho-joint-meeting-on-pesticide-specifications-jmps/pesticide-specifications/pesticide-specifications-list/en/>

4 Physical properties

4.1 pH range (MT 75.3, CIPAC Handbook J, p 131, 2000)

pH range: 4 to 7 (1% in water).

4.2 Dry sieve test (MT 59.1, CIPAC Handbook F, p. 177, 1995)

Maximum: 5% retained on a 75 µm test sieve.

Not more than $(0.005 \times X)\%$ of the mass of the sample used for the test shall be present as propoxur in the residue on the sieve, where X is the propoxur content (g/kg) found under 2.2 (Note 1).

5 Storage stability

5.1 Stability at elevated temperature (MT 46.4, CIPAC Handbook P, p. 232, 2021)

After storage at $54 \pm 2^\circ\text{C}$ for 14 days, the determined average active ingredient content must not be lower than 97% relative to the determined average content found before storage (Note 2) and the formulation shall continue to comply with the clauses for:

- pH range (4.1);
- dry sieve test (4.2).

Note 1 For example, if the formulation has a determined content of 40 g/kg of propoxur and 20 g of sample is used in the test, then the amount of propoxur in the residue on the sieve should not exceed 0.040 g, i.e.

$$\frac{(0.005 \times 40) \times 20}{100} \text{ g}$$

Note 2 Samples of the formulation taken before and after the storage stability test should be analyzed together after the test in order to reduce the analytical error.

PROPOXUR WETTABLE POWDER

FAO Specification 80 / WP (October 2021*)

This specification, which is PART ONE of this publication, is based on an evaluation of data submitted by the manufacturers whose names are listed in the evaluation reports (80/2003 & 80/2016). It should be applicable to relevant products of these manufacturers but it is not an endorsement of those products, nor a guarantee that they comply with the specification. The specification may not be appropriate for the products of other manufacturers. The evaluation reports (80/2003 & 80/2016), as PART TWO, form an integral part of this publication.

1 Description

The material shall consist of an homogeneous mixture of technical propoxur, complying with the requirements of FAO Specification 80/TC (October 2021). It shall be in the form of a fine, beige powder free from visible extraneous matter and hard lumps.

2 Active ingredient

2.1 Identity tests (80/TC/M2/2, CIPAC Handbook D, p.155, 1988)

The active ingredient shall comply with an identity test and, where the identity remains in doubt, shall comply with at least one additional test.

2.2 Propoxur content (80/WP/M2/3, CIPAC Handbook D, p.155, 1988)

The propoxur content shall be declared (g/kg) and, when determined, the average content measured shall not differ from that declared by more than the following tolerances.

Declared content in g/kg	Tolerance
above 100 up to 250	± 6% of the declared content
above 250 up to 500	± 5% of the declared content
Note: in each range the upper limit is included	

3 Relevant impurities

3.1 Water (MT 30.6, CIPAC Handbook P, p. 222, 2021)

Maximum: 20 g/kg.

* Specifications may be revised and/or additional evaluations may be undertaken. Ensure the use of current versions by checking at: <https://www.fao.org/pest-and-pesticide-management/guidelines-standards/faowho-joint-meeting-on-pesticide-specifications-jmps/pesticide-specifications/pesticide-specifications-list/en/>

4 Physical properties

4.1 pH range (MT 75.3, CIPAC Handbook J, p.131, 2000)

pH range: 4.5 to 7.5.

4.2 Wet sieve test (MT 185, CIPAC Handbook K, p.149, 2003)

Maximum: 2% retained on a 75 µm test sieve.

4.3 Suspensibility (MT 184.1, CIPAC Handbook P, p.245, 2021) (Note 1)

A minimum of 60% of the propoxur content found under 2.2 shall be in suspension after 30 min in CIPAC Standard Water D at $30 \pm 2^\circ\text{C}$ (Note 2).

4.4 Persistent foam (MT 47.3, CIPAC Handbook O, p.177, 2017) (Note 3)

Maximum: 10 ml after 1 min.

4.5 Wettability (MT 53.3.1, CIPAC Handbook F, p.164, 1995)

The formulation shall be completely wetted in 2 min without swirling.

5 Storage stability

5.1 Stability at elevated temperature (MT 46.4, CIPAC Handbook P, p. 232, 2021)

After storage at $54 \pm 2^\circ\text{C}$ for 14 days, the determined average active ingredient content must not be lower than 97% relative to the determined average content found before storage (Note 4) and the formulation shall continue to comply with the clauses for:

- pH range (4.1);
- wet sieve test (4.2);
- suspensibility (4.3);
- wettability (4.5).

Note 1 The formulation should be tested at the highest and lowest rates of use recommended by the supplier, provided this does not exceed the conditions given in method MT 184.1

Note 2 Chemical assay is the only fully reliable method to measure the mass of active ingredient still in suspension. However, simpler methods such as gravimetric and solvent extraction determination may be used on a routine basis provided that these methods have been shown to give equal results to those of chemical assay. In case of dispute, chemical assay shall be the "referee method".

Note 3 The mass of sample to be used in the test should correspond to the highest rate of use recommended by the supplier. The test is to be conducted in CIPAC standard water D.

Note 4 Samples of the formulation taken before and after the storage stability test may be analyzed concurrently after the test in order to reduce the analytical error.

PROPOXUR EMULSIFIABLE CONCENTRATE

FAO Specification 80 / EC (October 2021*)

This specification, which is PART ONE of this publication, is based on an evaluation of data submitted by the manufacturer whose name is listed in the evaluation report (80/2003). It should be applicable to relevant products of this manufacturer but it is not an endorsement of those products, nor a guarantee that they comply with the specification. The specification may not be appropriate for the products of other manufacturers. The evaluation report (80/2003), as PART TWO, forms an integral part of this publication.

1 Description

The material shall consist of technical propoxur, complying with the requirements of FAO Specification 80/TC (October 2021), dissolved in suitable solvents, together with any other necessary formulants. It shall be in the form of a stable, clear homogeneous, slightly yellow liquid, free from visible suspended matter and sediment, to be applied as an emulsion after dilution in water.

2 Active ingredient

2.1 Identity tests (80/TC/M2/2, CIPAC Handbook D, p. 155, 1988)

The active ingredient shall comply with an identity test and, where the identity remains in doubt, shall comply with at least one additional test.

2.2 Propoxur content (80/EC/M2/3, CIPAC Handbook D, p. 155, 1988)

The propoxur content shall be declared (g/kg or g/l at $20 \pm 2^\circ\text{C}$, Note 1) and, when determined, the average content measured shall not differ from that declared by more than the following tolerance.

Declared content in g/kg or g/l at $20 \pm 2^\circ\text{C}$	Tolerance
above 100 up to 250	$\pm 6\%$ of the declared content
Note: the upper limit is included in the range	

3 Relevant impurities

3.1 Water (MT 30.6, CIPAC Handbook P, p. 222, 2021)

Maximum: 10 g/kg.

* Specifications may be revised and/or additional evaluations may be undertaken. Ensure the use of current versions by checking at: <https://www.fao.org/pest-and-pesticide-management/guidelines-standards/faowho-joint-meeting-on-pesticide-specifications-jmps/pesticide-specifications/pesticide-specifications-list/en/>

4 Physical properties

4.1 **pH range** (MT 75.3, CIPAC Handbook J, p 131, 2000)

pH range: 3.0 to 4.2 (at 1% in water)

4.2 **Emulsion stability and re-emulsification** (MT 36.3, CIPAC Handbook K, p. 137, 2003) (Note 2)

The formulation, when diluted at $30 \pm 2^\circ\text{C}$ with CIPAC Standard Waters A and D, shall comply with the following:

Time after dilution	Limits of stability
0 h	Initial emulsification complete
0.5 h	"Cream", maximum: 1 ml
2.0 h	"Cream", maximum: 2 ml "Free oil", maximum: 0 ml
24 h	Re-emulsification complete
24.5 h	"Cream", maximum: 1 ml "Free oil", maximum: 0 ml
Note: in applying MT 36.3, tests after 24 h are required only where results at 2 h are in doubt	

5 Storage stability

5.1 **Stability at 0°C** (MT 39.3, CIPAC Handbook J, p.126, 2000)

After storage at $0 \pm 1^\circ\text{C}$ for 7 days, the volume of solid and/or liquid which separates shall not be more than 0.3 ml.

5.2 **Stability at elevated temperature** (MT 46.4, CIPAC Handbook P, p. 232, 2021)

After storage at $54 \pm 2^\circ\text{C}$ for 14 days, the determined average active ingredient content must not be lower than 97% relative to the determined average content found before storage (Note 3) and the formulation shall continue to comply with the clauses for:

- pH range (4.1);
- emulsion stability and re-emulsification (4.2).

Note 1 If the buyer requires both g/kg and g/l at 20°C , then in case of dispute the analytical results shall be calculated as g/kg.

Note 2 This test will normally only be carried out after the heat stability test, clause 5.2.

Note 3 Samples of the formulation taken before and after the storage stability test may be analyzed concurrently after the test in order to reduce the analytical error.

PART TWO

EVALUATION REPORTS

PROPOXUR

	Page
2021 FAO/WHO evaluation report based on submission of data from Hunan Haili Chemicals Limited (TC)	12
Supporting information	14
Annex 1: Hazard summary provided by the proposer	16
Annex 2: References	18
2016 FAO/WHO evaluation report based on submission of data from Tagros Chemicals India Limited (TC, WP, WP-SB)	19
Supporting information	23
Annex 1: Hazard summary provided by the proposer	25
Annex 2: References	27
2003 FAO/WHO evaluation report based on submission of data from Bayer CropScience (TC, DP, WP, EC)	28

PROPOXUR

FAO/WHO EVALUATION REPORT 80/2021

Recommendations

The Meeting recommended the following.

- (i) The propoxur TC produced by Hunan Haili Chemical Industry Co., Ltd. should be accepted as equivalent to the propoxur reference profile.
- (ii) The existing FAO specifications for propoxur TC should be extended to encompass the corresponding product of Hunan Haili Chemical Industry Co., Ltd.

Appraisal

The Meeting considered information, draft specification and data submitted by Hunan Haili Chemical Industry Co., Ltd. (Hunan Haili) in support of extension of the existing FAO specification for propoxur TC. The data submitted by Hunan Haili were broadly in accordance with the requirements of the Manual on development and use of FAO and WHO specifications for pesticides (2016 - 3rd revision of the 1st Edition).

Hunan Haili provided the Meeting with confidential information on the manufacturing process, 5-batch analysis data and manufacturing quality control limits for propoxur and all detectable impurities. The company stated that the confidential data submitted to JMPS are the same as those submitted for registration in Mexico and a valid registration certificate was also provided.

Hunan Haili utilize a manufacturing process for propoxur TC that differs from the reference process (Bayer CropScience), mainly for the starting materials used in the synthesis, but the overall route is similar.

The purity / impurity profile was supported by a GLP 5-batch analysis study. The batches of propoxur TC were manufactured in 2017.

The content of propoxur was determined by reverse phase HPLC, following the CIPAC method 80/TC/M2/3 published in CIPAC Handbook D.

The manufacturing impurities were determined by reverse phase HPLC, except for residual water which was determined using the CIPAC Karl Fischer method MT 30.5. Acetone insolubles and free acidity were also determined according to the CIPAC methods MT 27 and MT 31 respectively. All the analytical methods used for impurities were properly validated with regard to their specificity (with additional confirmation by HPLC-MS for propoxur related impurities), linearity of response, accuracy, precision and limits of detection (LOD) and quantification (LOQ).

The minimum purity of propoxur in the TC is 980 g/kg and complies with the existing FAO specification. Water and material insoluble in acetone are relevant impurities, and the control limits of these impurities are in line with the existing specification. Mass balances for the 5 batches are very high (99.0 - 99.2%), with no unknown detected, and similar to those of the reference profile of Bayer CropScience (99.9 - 100.1%). The manufacturing limit for a new impurity was reported at 1.2 g/kg originally based on the 5-batch analysis study. But the levels

determined in the batches used in the 5-batch analysis study were all below 1 g/kg and QC data showed that the found concentration levels of this impurity in 30 recent batches of propoxur TC were all below 1 g/kg.

The Meeting noted the inconsistency between the manufacturing limits of QC and the value reported. The company confirmed that their actual manufacturing limit was 1 g/kg and that 1.2 g/kg was calculated by mean value plus three times the standard deviations as the Manual suggests.

The Meeting requested Hunan Haili to provide a reasoned case, with data if available as to why the new impurity is not relevant. The company explained why the new impurity should be deemed relevant based on the criteria set out in the Manual. However the case was not fully accepted by the Meeting. Subsequently the company submitted an Ames test report, and the new impurity was determined to be negative for bacterial reverse mutation test. The proposer was questioned about the possible residue of an intermediate, and the proposer replied that the chemical compound was very volatile, and they had a process in place to remove its residue. QC data indicated the residual concentration level of the intermediate in 30 recent batches of propoxur TC were all below the LOQ of 0.08 g/kg. The water content, material insoluble in acetone and acidity measured in the 5 batches fully comply with the FAO/WHO specifications for propoxur TC.

An *in-vitro* reverse mutation assay with propoxur TC following the OECD guideline 471 was provided. The study was performed in compliance with GLP, and the results led to the conclusion that the propoxur TC from Hunan Haili does not induce reverse mutations under the conditions of the test.

No studies on physico-chemical properties of pure propoxur nor on the toxicology other than mutagenicity data nor ecotoxicology profiles were provided by Haili.

The Meeting concluded that based on the Tier-1 data provided by Hunan Haili that propoxur TC produced by Hunan Haili can be considered as equivalent to the reference profile supporting the existing FAO and WHO TC specifications (FAO/WHO evaluation report 80/2003).

Furthermore, the Meeting recommended to editorially update the propoxur specifications for TC, DP, WP and EC to bring them in line with the latest versions of the model specifications for these compounds. The adaptations introduced include e.g.

- The determination of residual water in the formulations now utilizes the harmonized method MT 30.6 (volumetric submethod of the Karl Fischer titration) published in Handbook P
- The determination of the suspensibility is now with the revised MT 184.1
- The accelerated storage test now references to the harmonized MT 46.4
- On the other hand, the emulsion stability test MT 36.3 used in the EC specification offering some discretion in the choice of temperature range where the test is carried out (e.g. 25 or 30 °C ± 2 °C, see Note 2) has been kept at 30 ± 2 °C, as the actual limits had been determined at 30 ° and the temperature dependence of the results are not yet fully explored for all EC formulations.

**SUPPORTING INFORMATION
FOR
EVALUATION REPORT 80/2021**

Table 1. Chemical composition and properties of propoxur technical material (TC)

Manufacturing process, maximum limits for impurities ≥ 1 g/kg, 5 batch analysis data	Confidential information supplied and held on file by FAO. Mass balances were 99.0 - 99.2% and percentages of unknowns were 0.80 - 0.97%.
Declared minimum propoxur content	980 g/kg
Relevant impurities ≥ 1 g/kg and maximum limits for them	Water, 1.6 g/kg Material insoluble in acetone, 0.6 g/kg
Relevant impurities < 1 g/kg and maximum limits for them	None
Stabilisers or other additives and maximum limits for them	None

Melting temperature range of the TC and/or TK	Crystal modification I: 87.2-88.7 °C Crystal modification II: 93.6-95.9 °C	98.6	EPA OCSP 830.7200, OECD 102, EEC A.1	ABC-2017-055
Solubility in organic solvents	n-Hexane:1.09 g/l at 20 °C Methanol:398.78 g/l at 20 °C	98.6	OCSP 830.7840, OECD 105, EEC A.6	ABC-2017-055

Methods of analysis and testing

Propoxur is determined using the CIPAC method by reverse phase HPLC with internal standardization using butyrophene and UV detection at 280 nm. Propoxur is identified by HPLC retention time and by IR and mass spectra.

The methods for determination of impurities were based on reverse phase HPLC.

Containers and packaging

Propoxur should be packed in polyethylene or polyamide using additional outer packaging.

ANNEX 1

HAZARD SUMMARY PROVIDED BY THE PROPOSER

Note:

Hunan Haili Chemical Industry Co., Ltd. provided written confirmation that the toxicological data included in the following summary were derived from propoxur having impurity profiles similar to those referred to in Table 1, above.

Table 2. Mutagenicity profile of deltamethrin technical material based on in vitro and in vivo tests

Species	Test	Purity %	Guideline, duration, doses and conditions	Result	Study number
<i>Salmonella typhimurium</i> , TA 98, TA 100, TA 1535, TA 1537, TA 102	<i>In vitro</i> genotoxicity testing - Bacterial reverse mutation test (Ames test)	98.5	Guideline: OECD No.471.1997, concentration 0.05, 0.16, 0.50, 1.6 and 5 mg/plate, in the absence and presence of S9	Non-mutagenic	BIO-GT 991

ANNEX 2: REFERENCES

Study number	Author(s)	Year	Study title. Study identification number. Report identification number. GLP [if GLP]. Company conducting the study
	FAO/WHO	2016	Manual on development and use of FAO and WHO specifications for pesticides. 2016 Third Revision of First Edition. FAO Plant Production and Protection Paper http://www.fao.org/fileadmin/templates/agphome/documents/Pests_Pesticides/Specs/JMPS_Manual_2016/3rd_Amendment_JMPS_Manual.pdf
ABC-2017-054	Yeming Zhang	2018	Five-Batch Analysis of Propoxur TGAI - Validation of Analytical Methodology for the Assay of Active Ingredient and Impurities in Propoxur TGAI and Subsequent 5-Batch Analysis of the Test Item. GLP. Achiever Biochem Co., Ltd. GLP Lab, P. R. China. Unpublished.
ABC-2017-055	Yeming Zhang	2018	Physical and Chemical Characterization of Propoxur TGAI - Color, Physical State, Odor, Stability and Corrosive Characteristics, Oxidation/Reduction: Chemical Incompatibility, pH, UV/Visible Absorption, Melting Point, Density, Partition coefficient, Solubility and Vapor Pressure. GLP. Achiever Biochem Co., Ltd. GLP Lab, P. R. China. Unpublished.
BIO-GT 991		2018	Bacterial Reverse Mutation Test of Propoxur Tech. using <i>Salmonella typhimurium</i> Tester Strains. GLP. Unpublished.
21056		2021	Bacterial Reverse Mutation Test with Propoxur Impurity 2 using <i>Salmonella typhimurium</i> . GLP. Unpublished.

PROPOXUR

FAO/WHO EVALUATION REPORT 80/2016

Recommendations

The Meeting recommended the following.

- (i) The propoxur TC as proposed by Tagros Chemicals India Limited should be accepted as equivalent to the propoxur reference profile.
- (ii) The existing FAO specifications for propoxur TC and WP should be extended to encompass the corresponding products of Tagros Chemicals India Limited.
- (iii) The existing WHO specifications for propoxur TC and WP should be extended to encompass the corresponding products of Tagros Chemicals India Limited.
- (iv) The specification for propoxur WP-SB, proposed by Tagros Chemicals India Limited, as amended, should be adopted by WHO.

Appraisal

The Meeting considered information, specifications and data submitted by Tagros Chemicals India Limited (India) in support of extension of the existing FAO and WHO specifications for propoxur TC and WP and in support of a new WHO specification for propoxur WP-SB. The data submitted by Tagros were broadly in accordance with the requirements of the Manual on development and use of FAO and WHO specifications for pesticides (November 2010 - second revision of the First Edition).

Propoxur TC

Tagros provided the Meeting with confidential information on the manufacturing process, 5-batch analysis data and manufacturing quality control limits for propoxur and all detectable impurities.

Tagros stated that their propoxur TC has been submitted for registration in Indonesia. The confidential information (manufacturing process, purity and impurity profile) submitted to FAO/WHO was confirmed by the Indonesian registration authorities as being identical to that submitted for registration in Indonesia, and was evaluated and considered acceptable by the Indonesian registration authorities.

The manufacturing process of propoxur TC from Tagros differs from the reference process (Bayer CropScience), mainly for one of the starting materials used in the synthesis and for the solvent used in the crystallization step, but the principle remains the same.

The purity / impurity profile was supported by a GLP 5-batch analysis study. The batches of propoxur TC were manufactured from September 2013 to January 2014.

Propoxur content was determined by reverse phase HPLC-DAD after dissolution in acetonitrile and internal standard calibration. The method was fully validated on its specificity (with additional confirmation by LC-MS), linearity of response, accuracy and precision. This method is similar to the CIPAC method 80/TC/M2/3 published in Handbook D, except that ethyl benzoate was used as internal standard instead of butyrophenone, the weight of propoxur in the calibration and samples solutions was decreased in order to avoid further dilution, and the chromatographic conditions were slightly adapted. The manufacturer explained that the CIPAC method for propoxur was developed in the 1980's and since that

time major advances in HPLC column technology have taken place. In the 5-batch analysis study, a number of potential internal standards were evaluated and of those, ethyl benzoate was found to be completely resolved from all of the known impurities of propoxur. The Meeting agreed that both methods are equivalent.

The propoxur manufacturing impurities and residual solvents were determined by reverse phase HPLC-DAD and GC-FID, except water which was determined using the CIPAC method MT 30.5. Acetone insolubles and free acidity were also determined according to the CIPAC methods MT 27 and MT 31 respectively. All the analytical methods used for impurities were fully validated on their specificity (with additional confirmation by LC-MS for propoxur related impurities), linearity of response, accuracy, precision and limits of detection (LOD) and quantification (LOQ).

The minimum purity of propoxur in the TC is 980 g/kg and complies with the existing FAO/WHO specification. No relevant impurities were declared. Mass balances for the 5 batches are high (98.7 - 99.6%), with no unknown detected, and similar to those of the reference profile of Bayer CropScience (99.9 - 100.1%). The 5-batch analysis study report indicates that no other significant impurity (each at or above 1 g/kg) was found in any of the 5 batches and that there is no indication that the assays employed missed any significant process related impurity. The water content, material insoluble in acetone and acidity measured in the 5 batches fully comply with the FAO/WHO specifications for propoxur TC.

The manufacturer was questioned about the possible presence of a chlorinated solvent in the final TC. The manufacturer provided a GLP 5-batch study showing that this chlorinated solvent measured by a GC-MS validated method was not detected in any of the 5 batches above the LOQ level of 0.05 g/kg, which is well below 10% of the GHS level of 1 g/kg.

Mutagenicity data were provided on *Salmonella typhimurium* (reverse mutation Ames test) following the OECD guideline 471. The study was performed in compliance with GLP, and the results led to the conclusion that the propoxur TC from Tagros is considered as non-mutagenic.

No studies on physico-chemical properties of pure propoxur nor on the toxicology and ecotoxicology profiles were provided by Tagros.

On basis of Tier-1 data provided by Tagros (manufacturing process, purity / impurity profile, 5-batch analysis data, mutagenicity profile), the Meeting concluded that the propoxur TC from Tagros should be considered as equivalent to the reference profile supporting the existing FAO and WHO specifications (FAO/WHO evaluation report 80/2003).

Propoxur WP and WP-SB

Tagros provided the Meeting with specifications for propoxur WP and WP-SB. The specification for the WP was similar to the existing FAO/WHO specifications for propoxur WP. The new WP-SB specification was supported by GLP studies performed on one single batch as well as quality control data on several batches of propoxur WP-SB.

Propoxur WP is recommended by WHOPEs for indoor residual spraying against malaria vectors at a dosage of 1-2 g a.i./m² for an effective action duration of 3-6 weeks (WHO 2006).

Tagros submitted GLP reports on appearance, propoxur content, water content, pH of a 10% aqueous suspension, wet sieve test, suspensibility, persistent foam, wettability, dissolution of the water soluble bag and accelerated storage stability at 54°C for 14 days. The analytical method used for propoxur content was the CIPAC method 80/WP/M2/3 published in CIPAC Handbook D and was fully validated on its specificity, linearity of response, accuracy and precision. The methods used for the physico-chemical tests were all CIPAC methods. Nevertheless, pH was performed in CIPAC water D instead of in distilled or de-ionized water, but it was considered acceptable by the Meeting. The manufacturer of the reference specification (Bayer CropScience) was questioned about the pH clause at 10% in water while the CIPAC method MT 75.3 recommends to perform the test at 1%. Bayer CropScience agreed to revise the pH range from 4.5 to 7.5% at 1% in water.

The suspensibility test was initially performed on the WP in presence of the soluble bag at a 1.25% a.i. concentration which is the maximum rate of use recommended by the supplier. As the suspensibility index can vary depending on the concentration of use of the product, the FAO/WHO specification guideline for WP and WP-SB recommends to perform the suspensibility test at the highest and lowest rates of use recommended by the supplier. At the request of the Meeting, the manufacturer provided additional suspensibility data at the minimum (1% a.i.) and maximum (1.25% a.i.) recommended concentrations with and without the soluble bag, and before and after accelerated storage.

The persistent foam test was performed on the WP with and without the soluble bag. Despite different concentrations were used in the test (4% a.i. for the WP and 1% a.i. for the WP-SB), these concentrations are higher or quite close to the highest rate of use of 1.25% a.i. recommended by the manufacturer, and it was considered acceptable by the Meeting.

Tagros initially specified a tolerance of 95% in the WP-SB specification for the active ingredient content remaining after accelerated storage at 54°C for 14 days while the tolerance is 97% in the existing FAO/WHO specification for propoxur WP. The results from the GLP study on propoxur WP-SB showed that the active ingredient after accelerated storage is well higher than 97% relative to the content before storage and that the product still complies with the clauses for pH range, wet sieve test, suspensibility, persistent foam, wettability and dissolution of the soluble bag. The manufacturer finally agreed to comply with the 97% tolerance for the active ingredient after accelerated storage.

The results from the GLP studies and the quality control data on the propoxur WP and WP-SB from Tagros showed that their WP fully comply with the existing FAO/WHO specification for the neat WP formulation and that their WP-SB fully comply with the proposed specification for the WP-SB.

The Meeting agreed also :

- to update in the specification for propoxur WP the CIPAC methods for wet sieve test (MT 185 instead of MT 59.3), suspensibility (MT 184 instead of MT 15.1 or MT 177) and persistent foam (MT 47.3 instead of MT 47.2);

- and to update in the specification for propoxur EC the CIPAC method for emulsion stability (MT 36.3 instead of MT 36.1.1)
to be in line with the current CIPAC methods.

**SUPPORTING INFORMATION
FOR
EVALUATION REPORT 80/2016**

Physico-chemical properties of propoxur

Table 1. Chemical composition and properties of propoxur technical material (TC)

Manufacturing process, maximum limits for impurities ≥ 1 g/kg, 5 batch analysis data	Confidential information supplied and held on file by FAO and WHO. Mass balances were 98.7-99.6 % with no unknowns.
Declared minimum propoxur content	980 g/kg
Relevant impurities ≥ 1 g/kg and maximum limits for them	None
Relevant impurities < 1 g/kg and maximum limits for them	None
Stabilisers or other additives and maximum limits for them	None

Methods of analysis and testing

Propoxur is determined by HPLC using internal standardization with ethyl benzoate and UV detection at 280 nm. This method is similar to the CIPAC method except that ethyl benzoate is used as internal standard instead of butyrophenone. Propoxur is identified by HPLC retention time and by IR and mass spectra.

The methods for determination of impurities were based on HPLC.

Containers and packaging

Propoxur should be packed in polyethylene or polyamide using additional outer packaging.

ANNEX 1

HAZARD SUMMARY PROVIDED BY THE PROPOSER

Note:

Tagros Chemicals India Limited provided written confirmation that the toxicological data included in the following summary were derived from propoxur having impurity profiles similar to those referred to in Table 1, above.

All data has been generated only with the Tagros technical grade active ingredient.

Table A. Mutagenicity profile of propoxur technical material based on *in vitro* tests

Species	Test	Purity %	Guideline, duration, doses and conditions	Result	Study number
<i>Salmonella typhimurium</i> (TA1537, TA1535, TA102, TA100 and TA98)	Reverse Mutation Assay (Ames test), <i>in vitro</i>	98.20%	OECD No. 471 Dosage: 128, 320, 800, 2000 and 5000 µg/plate Solvent: DMSO	Negative	14_14_058

ANNEX 2: REFERENCES

Study number	Author(s)	Year	Study title. Study identification number. Report identification number. GLP [if GLP]. Company conducting the study
EPP00146	W Bruce Craig	2014	5-Batch Analysis of Propoxur TGAI in Accordance with Regulation (EC) No. 1107/2009, Referencing SANCO 3030/99 rev. 4. Study No. AN140129A. Report No. EPP00146. EPP Limited, UK. GLP, Unpublished.
14_14_058	Kapil Nikam	2014	Bacterial Reverse Mutation Test of Propoxur Technical in Salmonella typhimurium Tester Strains. Study No. 14_14_058. Sa-FORD, India. GLP, Unpublished.
5083	Nageswara T.	2015	Determination of Foam Persistence of Propoxur 50% WP-SB. Study No. 5083. RCC Laboratories India Private Limited, India. GLP, Unpublished.
5084	Nageswara T.	2015	Determination of Accelerated Storage Stability and Relevant Physico-chemical properties (Foam Persistence) of Propoxur 50% WP-SB. Study No. 5084. RCC Laboratories India Private Limited, India. GLP, Unpublished.
G9778	Ravikanth Gogineni	2014	Method validation and active ingredient analysis of propoxur 50% WP-SB. Study No. G9778. Advinius Therapeutics Limited, India. GLP, Unpublished.
G9779	Shanthaveer appa K. S.	2014	Determination of color, odor and physical state of propoxur 50% WP-SB. Study No. G9779. Advinius Therapeutics Limited, India. GLP, Unpublished.
G9780	Shanthaveer appa K. S.	2014	Determination of moisture content of propoxur 50% WP-SB. Study No. G9780. Advinius Therapeutics Limited, India. GLP, Unpublished.
G9781	Shanthaveer appa K. S.	2014	Determination of pH of 10% (w/v) aqueous suspension of propoxur 50% WP-SB. Study No. G9781. Advinius Therapeutics Limited, India. GLP, Unpublished.
G9782	Shanthaveer appa K. S.	2014	Wet sieve test for propoxur 50% WP-SB. Study No. G9782. Advinius Therapeutics Limited, India. GLP, Unpublished.
G9783	Shanthaveer appa K. S.	2014	Determination of suspensibility of propoxur 50% WP-SB. Study No. G9783. Advinius Therapeutics Limited, India. GLP, Unpublished.
G9784	Shanthaveer appa K. S.	2014	Determination of persistent foam of propoxur 50% WP-SB. Study No. G9784. Advinius Therapeutics Limited, India. GLP, Unpublished.
G9785	Shanthaveer appa K. S.	2014	Determination of wettability of propoxur 50% WP-SB. Study No. G9785. Advinius Therapeutics Limited, India. GLP, Unpublished.
G9786	Shanthaveer appa K. S.	2014	Determination of dissolution rate of water soluble bags of propoxur 50% WP-SB. Study No. G9786. Advinius Therapeutics Limited, India. GLP, Unpublished.
G9787	Shanthaveer appa K. S.	2014	Accelerated storage stability test of propoxur 50% WP-SB. Study No. G9787. Advinius Therapeutics Limited, India. GLP, Unpublished.
DNA3080	Norris David	2015	Analysis of 5 batches of Propoxur Technical to determine the content of a specified solvent, with associated validation, in Compliance with Good Laboratory Practice. Study Number DNA3080. David Norris Laboratories Ltd. GLP, Unpublished.
	WHO	2006	Pesticides and their application for the control of vectors and pests of public health importance (sixth edition). Document WHO/CDS/NTD/WHOPES/GCDPP/2006.1. WHO, 2006. Available at http://whqlibdoc.who.int/hq/2006/WHO_CDS_NTD_WHOPES_GCDPP_2006.1_eng.pdf?ua=1

PROPOXUR

FAO/WHO EVALUATION REPORT 80/2003

Explanation

The data for propoxur were evaluated for review of existing FAO and WHO specifications for propoxur. Existing FAO full specifications for TC, DP, WP and EC (80/TC/S/F, 1991; 80/DP/S/F, 1992; 80/WP/S/F, 1991; 80/EC/S/F, 1991) were published as AGP: CP/332 in 1995. Existing WHO full specifications for TC and WP (WHO/SIT/18.R4 and WHO/SIF/30.R4, respectively) were published in 1999.

Propoxur is no longer under patent.

Propoxur was evaluated by the FAO/WHO JMPR in 1973, 1977, 1981, 1983 1989 and 1996. Propoxur has also been submitted to the EU as notification for the Biocidal Products Directive (Directive 98/8/EC) under notification no. N353.

The draft specification and the supporting data were provided by Bayer Crop Science AG, Germany, in 2002.

Uses

Propoxur is an *N*-methylcarbamate insecticide and acaricide. It is non-systemic is a contact and stomach poison, which does not accumulate. The mode of action is interference with nervous transmission across the synaptic gap through inhibition of acetylcholinesterase.

Propoxur is used both for agricultural and public health purposes, being applied by spraying or as a dust. It is used against insect pests such as chewing and sucking insects, ants, cockroaches, crickets, flies and mosquitoes. Agricultural crop applications include sugar cane, cocoa, grapes and other fruit, maize, rice, vegetables, cotton, lucerne, forestry and ornamentals.

Identity

ISO common name:

propoxur (E-ISO, F-ISO 1750)

Chemical names:

IUPAC: 2-isopropoxyphenyl methylcarbamate

CA: 2-(1-methylethoxy)phenyl methylcarbamate

CAS No:

114-26-1


CIPAC No:

80

Synonyms:


Bay 9010, Baygon, Bayer 39007, Blattanex, Bolfo, BO Q 5812315, OMS 33, PHC (JMAF), Pillargon, UN Carbamate, Tugon, Uden, Undene

Structural formula:



Solid propoxur can exist in two crystal forms (modifications I and II) but the technical material usually contains >95% of modification I.

Molecular formula:



Relative molecular mass:

209.25

Identity tests:

HPLC retention time, with detection at 280 nm (CIPAC Handbook D, p. 155, 1988); IR and mass spectra; melting point (87.5-90°C).

Physico-chemical properties of propoxur

Table 1. Physico-chemical properties of pure propoxur

Parameter	Value(s) and conditions	Purity %	Method reference
Vapour pressure:	1.29 mPa at 20°C 2.78 mPa at 25°C (extrapolated)	99.9	OECD 104
Melting point and temperature of decomposition:	Melting point: Crystal modification I: 87.5°C Crystal modification II: 90.0°C Stable under ambient conditions, no decomposition occurs below 150°C Decomposition starts at about 220°C and the consequent multi-stage process evolves about 300 kJ/kg.	99.9	OECD 102/113
Solubility in water:	1.75 g/l at 20°C	99.8	US-EPA Guidelines
Octanol/water partition coefficient:	log P _{ow} = 1.56 at 20°C	99.8	US-EPA Guidelines
Hydrolysis characteristics:	Half-life at 22°C >1 year at pH 4 93.2 days at pH 7 30.1 hours at pH 9	Not reported	OECD 111
Dissociation characteristics:	Propoxur has no distinct acidic or basic properties in aqueous solution.	99.9	OECD 112
Density	1.17 g/cm ³ at 20°C	99.9	OECD 109

Table 2. Chemical composition and properties of propoxur technical materials (TC)

Manufacturing process, maximum limits for impurities ≥ 1 g/kg, 5 batch analysis data.	Confidential information supplied and held on file by WHO. Mass balances were 99.9 to 100.1% with no unidentified impurities.
Declared minimum propoxur content:	980 g/kg
Crystal modification ratio of propoxur (discernible by IR)	Modification I: > 950 g/kg Modification II: < 50 g/kg
Relevant impurities ≥ 1 g/kg and maximum limits for them:	None
Relevant impurities < 1 g/kg and maximum limits for them:	None
Stabilisers or other additives and maximum limits for them:	None
Melting or boiling temperature range	87.5 to 90°C
Density	1.17 g/cm ³ at 20°C
Bulk density	0.52 kg/l

Hazard summary

Notes

(i) The Proposer confirmed that the toxicological and ecotoxicological data included in the summary below were derived from propoxur having impurity profiles similar to those referred to in the table above.

(ii) The conclusions expressed in the summary below are those of the proposer unless otherwise specified.

Table 3. Toxicology profile of propoxur technical material, based on acute toxicity, irritation and sensitization.

Species	Test	Duration and conditions	Purity	Result	Reference
Rat, Wistar, male, female	oral	60 animals, 14 days observation period, in polyethylene glycol 400, doses from 0 to 127.9 mg/kg bw	not stated	LD ₅₀ = Males: 89.7 mg/kg bw Females: 78.5 mg/kg bw	Sturdivant & Halliburton 1998
Rat, Wistar, male, female	oral	40 animals, in Lutrol, doses from 10 to 500 mg/kg bw	not stated	LD ₅₀ = Males: 196 mg/kg bw Females: 126 mg/kg bw	JMPR 1989, USEPA 1997, Flucke 1980
Rat, male, female	dermal	24 h, in Lutrol, applied to intact dorsal skin, observation period 14 days	99.6%	LD ₅₀ > 5000 mg/kg bw	JMPR 1989, USEPA 1997, Flucke 1980
Rat, male, female	inhalation	4 h, 40 animals, conc. in air from 0 to 912 mg/m ³	not stated	LC ₅₀ = 654 mg/m ³	Pauluhn 1993
Rat, male, female	inhalation	4 h, 5 animals per sex, con. 28.7, 110.1, 330.4, 497.5 mg/m ³	99.6%	LC ₅₀ > 0.5 mg/l	JMPR 1989, USEPA 1997, Pauluhn 1988
Rabbit	skin irritation	500 mg, 4 hours, 6 animals	not stated	No manifestations of irritation	USEPA 1997 Sheets & Fuss 1991

Species	Test	Duration and conditions	Purity	Result	Reference
Rabbit, New Zealand White	skin irritation	Dose not recorded 24 or 72 hours	99.2%	No manifestations of irritation	JMPR 1989 Thyssen 1978
Rabbit, New Zealand White	eye irritation	0.1 g, 9 animals, examinations 1, 24, 48, 72 and 96 h	99.6%	No manifestations of irritation Severe miosis, which disappeared within 24 hours, no signs of irritation up to 96 hours post-application	JMPR 1989 Yamane 1986b
Rabbit, New Zealand White	Eye irritation	65 mg, 6 males, 48 h	99.8%	Instillation resulted in minor eye irritation (redness and discharge) which cleared within 48 h	USEPA 1997 Sheets, 1990a
Guinea-pig	skin sensitization	Magnusson and Kligman test, 30 animals	98.8%	No evidence of skin-sensitizing potential	JMPR 1989 Heimann 1982a
Wistar rat, male, female	acute neurotoxicity	14 days, groups of 12 male and female rats, doses of 0, 2, 10, 25mg/kg	99.4%	NOEL could not be determined, LOEL = 2mg/kg, based on brain CHE inhibition in both sexes 45 min after dosing	USEPA 1997 Dreist & Popp 1994

Table 4. Toxicology profile of propoxur technical material based on repeated administration (sub-acute to chronic)

Species	Test	Duration and conditions	Purity	Result	Reference
Rat, Wistar male, female	oral, gavage	5 days, 5 animals per sex and , doses 0, 15, 30 mg/kg bw/day of propoxur of two different purities	98.6% technical and 99.2% recrystallized	Dose-related convulsions and apathy the only adverse effects, no difference between the two purities.	JMPR 1989 Heimann 1983
Wistar rat, female	Oral, feeding, toxicity	14 weeks, 100 rats, doses of 0, 8000 ppm via the feed	99.9%	NOAEL >8000 ppm	JMPR 1989 Hahnemann & Rühl-Fehlert 1988d
Wistar rat, female	Oral, feeding, toxicity	104 weeks, 610 animals, doses from 0, 50, 250, 1000, 3000, 5000, 8000 ppm (0 to 348.46 mg/kg bw/day)	99.6-99.9%	NOAEL = 250 ppm (14.47 mg/kg bw/day) Growth retardation, ChE inhibition, urinary bladder alterations. Hyperplastic and neoplastic changes to bladder were diet-dependent and hyperplastic changes were reduced by administration of ammonium chloride	JMPR 1989 Hahnemann & Rühl-Fehlert 1988d

Species	Test	Duration and conditions	Purity	Result	Reference
Mouse	Oral, feeding, toxicity	53 weeks, 50 female NMRI mice, diets containing 0, 3000, 8000 ppm	99.6-99.9%	Growth slightly decreased at 8000 ppm. Increased liver weight and fatty degeneration at 3000 and 8000 ppm. Relative lung weight increased at 8000 ppm only. No adverse effect on urinary bladder epithelium.	JMPR 1989 Hahnemann & Rühl-Fehlert 1988c
Hamster	Oral, feeding, toxicity	53 weeks, 50 female Syrian golden hamsters, diets containing 0, 3000, 8000 ppm.	99.6-99.9%	At both dose levels mortality incidence slightly increased, general state of animals impaired, growth retarded. Relative weights of kidneys and adrenals increased at 8000 ppm only. No adverse effect on urinary bladder epithelium.	JMPR 1989 Hahnemann & Rühl-Fehlert 1988a
Dog	Oral, feeding, toxicity	52 weeks, 12 Beagle dogs (m/f), 0, 200, 600 ppm. Additional groups weeks 1-40, 1800 ppm; weeks 41-44, 3600 ppm; weeks 45-52, 5400 ppm	99.4%	Cholinergic symptoms observed at highest dose level, after elevation of dose to 5400 ppm and 1/6 animals died. The following were also increased in this group: thrombocyte, leucocyte and reticulocyte counts, incidence of Heinz bodies, ALAT and SAP, liver weight and thyroid weight; thymus weight decreased. At highest dose and at 600 ppm, growth was retarded and plasma cholesterol and liver <i>N</i> -demethylase increased. NOAEL = 200 ppm.	JMPR 1989 USEPA 1997 Hoffmann & Gröning, 1984
Rhesus monkey	Oral, feeding, (intubation), toxicity	13 weeks, 6 rhesus monkeys (m/f), doses of 40 mg/kg bw/day	99.6%	Cholinergic symptoms observed but no adverse effect on urinary bladder epithelium.	JMPR 1989 USEPA 1997 Hoffmann & Rühl, 1985

Species	Test	Duration and conditions	Purity	Result	Reference
Mouse, male, female	Oral, feeding, carcinogenicity	SPF mice, strain CF1/W74, 2 years, groups of 50 male and 50 female rats, doses of 0, 700, 2000 or 6000 ppm in feed	99.6%	NOAEL = Males: 2000 ppm Females: 6000 ppm. No indications of oncogenic effects in any treatment group.	JMPR 1989 Bomhard & Löser 1981, Patterson 1980
Chinchilla rabbit, male, female	Dermal	5 male and 5 female rabbits, 14 applications, exposure period 24 hours, doses of 0 and 500 mg/kg bw/day	Not stated	Clinical, clinical chemical and haematological examinations did not detect any indications of damage or local irritant effects.	Kimmerle & Solmecke 1971
New Zealand white rabbit, male, female	Dermal	10 male, 10 female rabbits, 13 weeks, exposure period 6 hours/day, 5 days/week, doses of 0, 50, 250 and 1000 mg/kg bw/day in Cremophor	100%	NOAEL = 1000 mg/kg bw	USEPA 1997 Diesing & Flucke 1989
Wistar rat, male, female	inhalation	12 weeks, (6 h per day, 5 days a week), groups of 10 male and 10 female rats, concentrations of 0, 5.7, 18.7 or 31.7 mg/m ³	98.9%	Only effect observed was depression of cholinesterase activity in plasma, erythrocytes and brain, at 31.7 mg/m ³ only.	JMPR 1989, Kimmerle & Iyatomi 1976
Wistar rat, male, female	feeding, 2 generation reproduction	groups of 25 male and 25 female rats, duration 330 days, pre-mating exposure 70 days both groups, 0, 30 and 80 ppm in feed for entire period (preparation, mating, gestation and rearing)	99.8%	NOAEL = Parent: 30 ppm Reproduction: 80 ppm	USEPA 1997, Suter 1990, Dotti 1992

Species	Test	Duration and conditions	Purity	Result	Reference
Wistar rats, female	feeding, teratogenicity and embryotoxicity	25 mated females per group, exposure period from day 6 through 15 of gestation, in daily oral doses of 0, 3, 9 and 27 mg/kg bw, formulated in water/Cremophor EL	99.4%	NOAEL = 3 mg/kg bw/day for maternal toxicity No evidence of embryotoxicity or teratogenicity even at the highest dose tested (27 mg/kg bw).	JMPR 1989 USEPA 1997 Becker <i>et al.</i> 1989b
Rabbit, female, Chinchilla strain	feeding, teratogenicity and embryotoxicity	4 groups, 16 females per group, from 6 th to 18 th day of gestation, in daily oral doses of 0, 3, 10 and 30 mg/kg bw, formulation agent water/Cremophor EL	99.4%	NOAEL = 10 mg/kg bw/day for maternal toxicity and Embryotoxicity. Increased post-implantation losses at 30 mg/kg bw/day. Not teratogenic	JMPR 1989 USEPA 1997 Becker <i>et al.</i> 1989a
Wistar rat, male, female	sub-chronic neurotoxicity	13 weeks, groups of 12 male and 12 female rats, doses of 0, 500, 2000 and 8000 ppm equivalent to 0, 39, 163 and 703 mg/kg bw/day for females and 0, 33, 132 and 543 mg/kg bw/day males	99.5%	NOEL (functional observation battery and motor and locomotor activity changes) = males: 543 mg/kg bw Females: 163 mg/kg bw	USEPA 1997, Dreist & Popp 1994
White leghorn hens	sub-chronic delayed neurotoxicity	8 hens, 30 days, doses of 0, 300, 1500, 3000 and 4500 ppm	Not stated	No evidence of delayed neurotoxicity during feeding or 4 weeks post-treatment.	Kimmerle 1966a Hobik 1967
B6C3F1 mice	Oncogenicity	2 groups of 50 males and 50 females in 0, 500, 2000, 8000 ppm, 2 years	99.6%	NOEL = 500 ppm, LOEL = 2000 ppm	USEPA 1997 Bomhard 1992

Table 5. Mutagenicity profile of propoxur technical material based on *in vitro* and *in vivo* tests

Species	Test	Conditions	Purity	Result	Reference
<i>Salmonella typhimurium</i> (TA 100, TA 98, TA1535, TA 1537, TA 1538)	Ames test, <i>in vitro</i>	Concentrations: 50 nmol/plate	95%	Negative	JMPR 1989, Blevins <i>et al.</i> 1977b
<i>Salmonella typhimurium</i> (TA 100, TA 98, TA1535, TA 1537, TA 1538)	Ames test, <i>in vitro</i>	Concentrations: 0.1-1000µg/plate, solvent DMSO	98%	Negative	JMPR 1989, Inukai & Iyatomi 1978
<i>Escherichia coli</i> (WP2 hcr. B/r try WP2)	Reverse mutation test, <i>in vitro</i>	Concentrations: 20 µl/disk	Not stated	Negative	JMPR 1989 Shirasu <i>et al.</i> 1976
<i>Saccharomyces cerevisiae</i> (D4)	Mitotic gene conversion test, <i>in vitro</i>	Concentrations: 2 ml of suspension (containing 1000 ppm a.i.) at 5 x 10 cells; solvent DMSO	99.8%	Negative	JMPR 1989, Siebert & Lemperle 1974, Siebert & Eisenbrand 1974
Male mice	Dominant lethal test, <i>in vivo</i>	Concentrations: 10 mg/kg bw; p.o.	99.2%	Negative	JMPR 1989 Herbold 1980a
Male and female NMRI-mice bone marrow cells	Micronucleus test, <i>in vivo</i>	2 x 5 mg/kg bw; 2 x 10 mg/kg bw; p.o.	99.2%	Negative	JMPR 1989 Herbold 1980b

Table 6. Ecotoxicology profile of propoxur technical material

Species	Test	Duration and conditions	Purity	Result	Reference
<i>Daphnia magna</i> (water flea)	Acute toxicity	48 h	98.8%	EC ₅₀ = 0.011 ppm	USEPA 1997 Lamb 1981
<i>Lepomis macrochirus</i> (bluegill sunfish)	Short-term toxicity, flow-through	96h, concentrations from 2.2 to 10 ppm, temperature 22°C	98.8%	LC ₅₀ = 6.2 mg/l	USEPA 1997 Lamb 1981
Rainbow trout	Short-term toxicity, flow-through	96h, 5 concentrations: 2.2 to 10 ppm, temperature 22°C	98.8%	LC ₅₀ = 3.7 mg/l	USEPA 1997 Lamb 1981
<i>Scenedesmus subspicatus</i> (green algae)	Effect on growth, static water	72 h, Directive 92/69/EEC	99.6%	IC ₅₀ = 22 mg/l NOEC = 3.1 mg/l	Caspers 2001
Bobwhite quail	sub-acute toxicity	5 days, 10 birds per dietary level, doses of 500, 1000, 2000, 4000 and 8000 ppm	98.8%	LC ₅₀ = 2828 ppm NOEL = 1000 ppm	USEPA 1997 Lamb 1981

Propoxur was evaluated by WHOPES in 1976 and re-evaluated in 1999.

Propoxur was evaluated by the FAO/WHO JMPR in 1973, 1977, 1981, 1983, 1989, 1991 and 1996. with the toxicological reviews conducted in 1973 and 1989. The JMPR (JMPR 1989) concluded that propoxur showed moderate acute toxicity in the animal species examined. After reviewing all available data from *in vitro* and *in vivo* short-term tests, the JMPR concluded that there was no evidence of genotoxicity. The JMPR recommended an ADI of 0.02 mg/kg bw/day for propoxur.

The US EPA also evaluated propoxur (USEPA 1997) and concluded that it is likely to be moderately persistent under aerobic or anaerobic soil conditions (a metabolic half-life of several months), mobile (K_d values less than 1) and may potentially leach to groundwater. It is hydrolytically stable at acid or neutral pH (3-7) but degrades rapidly in alkaline conditions. Propoxur was categorized as very highly toxic to birds on an acute basis (some LD_{50} s are <10 mg/kg); highly toxic to birds on a sub-acute dietary basis (LC_{50} in the range of 51-500 ppm); moderately toxic to freshwater fish (some LC_{50} s in the range >1-10 ppm); highly toxic to bees (<11 µg/bee) on an acute contact basis; and very highly toxic to freshwater invertebrates (daphnid EC_{50} <1 ppm).

The WHO hazard classification of propoxur is: “moderately hazardous, class II” (WHO 2000) and the USEPA classification of acute toxicity is also class II (USEPA 1997).

Formulations

The main formulation types available are WP, DP and EC, which are registered and sold in more than 45 countries throughout the world.

Methods of analysis and testing

The analytical method for the active ingredient (which also provides an identity test) is CIPAC 80/TC/M/2/3. Propoxur is determined by HPLC, using internal standardization with butyrophenone and UV detection at 280 nm. Propoxur may be identified by HPLC retention time and by IR and mass spectra.

The methods for determination of impurities were based on HPLC.

Test methods for physico-chemical properties of technical active ingredient are OECD, EPA and EU, while those for the formulations are CIPAC, as indicated in the specifications.

Physical properties

The properties and limits proposed for the specifications for TC, WP and EC comply with the requirements of the WHO/FAO Manual (FAO/WHO 2002).

Containers and packaging

Propoxur should be packed in polyethylene or polyamide using additional outer packaging.

Expression of the active ingredient

The active ingredient is expressed as propoxur.

Appraisal

The Meeting considered data, provided by Bayer Crop Science AG, for the review of existing full FAO (TC, DP, WP, EC) and WHO (TC, WP) specifications for propoxur. Propoxur is no longer under patent, it is presently registered in more than 45 countries and

has been used in agriculture and public health applications for many years. It is, however, not approved for use in agriculture in the USA and the proposer reported that registration of propoxur for crop applications in Europe would not be supported.

Propoxur has been registered for many years in numerous countries world-wide. Information including that related to toxicology and ecotoxicology on propoxur is available from publications/websites of the US EPA, JMPR, WHO and EXTTOXNET (<http://exttoxnet.orst.edu/pips/propoxur.htm>). The Proposer stated that the data provided for this evaluation were similar to those provided to the JMPR for evaluation but was unable to state categorically that they were similar to those submitted to the US EPA (USEPA 1997).

Propoxur is an *N*-methyl carbamate insecticide which is fairly soluble in water, very soluble in polar organic solvents but only slightly soluble in non-polar organic solvents. It is hydrolyzed very slowly at pH 4, slowly at pH 7 but rather rapidly at pH 9.

Propoxur is of moderate mammalian toxicity, it is rapidly metabolized and does not accumulate in tissues. It is not sensitizing or irritant to skin and is not irritant to the eye, although transient severe miosis occurred following application to the eye. There is no evidence that propoxur is carcinogenic, teratogenic or embryotoxic (post-implantation loss occurred only at doses above the level at which maternal toxicity occurred). In a 5-day study on rats, comparing the toxicity of technical (purity 98.6%) and recrystallized (purity 99.2%) propoxur, no difference in toxicity was found. The JMPR has recommended an ADI of 0.02 mg/kg bw/day for propoxur.

As may be expected for such a carbamate insecticide, propoxur is highly toxic to honeybees, aquatic invertebrates and birds, though its toxicity varies according to the species. It is moderately to slightly toxic to fish. The reported 96-hour LC50 values are 3.7 mg/L in rainbow trout, and 6.6 mg/L in bluegill sunfish. Propoxur is highly toxic to freshwater invertebrates and very highly or highly toxic to birds, its toxicity varying according to species. Propoxur is rather persistent and mobile in soils, having characteristics which could produce leaching to groundwater.

The Meeting was provided with confidential information on the current manufacturing process, together with data from 5-batch analyses and the manufacturing specifications for all impurities ≥ 1 g/kg. Mass balances were high (99.9-100.1%) and no unidentified impurities were present. The current (2000-on) manufacturing process produces a higher purity TC than previously and no new impurities are found (Riegner 2005). The data were stated by the manufacturer to be identical to those submitted for registration in Mexico, Australia, the Philippines, Thailand, Venezuela and Malaysia. The data were confirmed as being essentially similar to those submitted to Australia (Sethi 2005).

The proposed specification for propoxur TC was in accordance with the requirements of the manual (FAO/WHO 2002), with the exception of the three clauses considered below. Meeting noted the proposed higher minimum purity of 980 g/kg (1991 FAO specification minimum 970 g/kg, 1999 WHO specification minimum 950 g/kg).

- (i) The manufacturer initially proposed a clause to control the crystal form ratio of propoxur TC, on the basis that crystal modification II has an adverse effect on the suspensibility of water dispersible formulations (Grohs 2004a). The two forms can apparently be distinguished by IR or x-ray diffraction methods but it was subsequently stated that the problem occurs only in WPs formulated with high concentrations (above 50%) of propoxur, which are no longer marketed (Grohs 2004b). The clause is not required for the low concentration and liquid formulations currently marketed and therefore the proposal was withdrawn.

(ii) A proposed clause for melting point of the TC, with a range of 87.5-90°C (instead of 86-91.5°C in the existing FAO and WHO specifications) was not in accordance with current guidelines in the manual (FAO/WHO 2002). The Meeting agreed that it should not be included but that it could be used as a supporting identity test.

(iii) The manufacturer explained (Grohs 2004a) that a proposed clause to limit acidity in the TC was necessary because, although propoxur is stable to hydrolysis in acid conditions, the TC is used to formulate water-based aerosols and the presence of excessive acid could initiate rapid rusting of the aerosol canister.

The proposed specifications for DP, WP and EC were broadly in accordance with the requirements of the manual (FAO/WHO 2002) but the following points were discussed and agreed with the manufacturer.

(i) The Meeting noted that the proposed minimum active ingredient content after the test of storage at elevated temperature was 95% (relative to the determined average content found before storage) compared to 97% in the existing FAO specification but the manufacturer subsequently confirmed that the limits should be 97% (Grohs 2004a).

(ii) The manufacturer initially specified the use of 63 µm sieve in the dry sieve test for the DP but agreed that the clause should be restricted to the standard 75 µm sieve (Grohs 2004a).

(iii) The manufacturer initially specified the use of a 40 µm sieve in the wet sieve test for the WP but agreed that the clause should be restricted to the standard 75 µm sieve (Grohs 2004a).

(iv) The Meeting questioned the long (2 min) wettability time specified for the WP. The manufacturer stated that propoxur is non-polar and therefore difficult to wet, that the 2 min limit is given in existing FAO and WHO specifications, and that the wettability had not given rise to practical problems in the field after many years of use (Grohs 2004a). Although it was noted that propoxur is not of exceptionally low polarity (it is slightly soluble in water), the Meeting agreed to accept the 2 min limit.

(v) The Meeting questioned the relatively high limit for water (10 g/kg) in the EC. The manufacturer stated that the EC is not turbid at <10 g/kg (Grohs 2004a) but that the water content must be kept below 10 g/kg in order to meet cold stability requirements. At temperatures below -5°C, ice crystals function as crystallization points and reduce the solubility of propoxur in the EC, causing sedimentation (Grohs 2004a). The Meeting noted that, although the proposed specification for propoxur WP (at the 500 g/kg level) is the same for both agricultural and public health applications, users should adhere to the label recommendations and not use the products interchangeably.

The analytical and physical test methods to be used in support of the proposed specifications are all CIPAC methods.

Recommendations

The Meeting recommended that:

- (i) the existing FAO specifications for propoxur TC, DP, WP and EC and the existing WHO specifications for propoxur TC and WP should be withdrawn;
- (ii) the proposed specifications (amended as described in the appraisal, above) for propoxur TC and WP should be adopted by FAO and WHO;

(iii) the proposed specifications (amended as described in the appraisal, above) for propoxur DP and EC should be adopted by FAO.

References

- Becker *et al.* 1989a Becker H., Mladenovic, P. & Terrier Ch. (1989a) Embryotoxicity study (including teratogenicity) with BOQ 5812315 (c.n. propoxur) in the rat. Report No. R 4686 (project 207270) dated March 1, 1989 from RCC-Research & Consulting Company Itingen, Switzerland, Unpublished.
- Becker *et al.* 1989b Becker H., Mladenovic, P. & Terrier Ch. (1989b) Embryotoxicity study (including teratogenicity) with BOQ 5812315 (c.n. propoxur) in the rabbit. Report No. R 4684 (Project 207292) dated March 1, 1989 from RCC-Research & Consulting Company Itingen, Switzerland, Unpublished.
- Blevins *et al.* 1977b Blevins, R.D., Lee, M. & Regan, J.D. (1977b) Mutagenicity screening of five methyl carbamate insecticides and their nitroso derivatives using mutants of *Salmonella typhimurium* LT2, *Mutat. Res.* 56: 1-6.
- Bomhard & Löser 1981 Bomhard E. and Löser E., (1981) B658 12 315 Chronic toxicological study on mice (Feeding study over two years). Report No. 9954 dated May 12, 1981 from Bayer A.G. Institute of Toxicology, Wuppertal-Elberfeld, Unpublished.
- Bomhard, 1992 Bomhard, E. (1992) BOQ 5812315 - Study for carcinogenicity in B6C3F1 mice (twenty-four month feeding study). Report No. 21571 dated July 27, 1992 from Bayer AG, Institute of Toxicology, Wuppertal.
- Caspers 2001 Caspers, N. (2001) BOQ 5812315 (propoxur) - Alga, growth inhibition test. Report No. 1071 A/01 AL, dated February 23, 2001, Bayer AG, Leverkusen.
- Diesing & Flucke 1989 Diesing, L. & Flucke, W. (1989) Subchronic Dermal Toxicity Study in Rabbits: BOQ 5812315 (Propoxur): Report No. 98568 (Mobay) prepared by Bayer AG, Unpublished.
- Dotti 1992 Dotti, A., Kinder, J., Biedermann, K., Luetkemeier, H. & Wright, J.; (1992) Two-generation reproduction study in rats. BOQ 5812315, Lab Project Number: 286997: T 5039706. Prepared by Research & Consulting Co. Ltd. And Biological Research Labs Ltd., Unpublished.
- Dreist & Popp 1994 Dreist, M., Popp, A. (1994) BOQ 5812315 (c.n. Propoxur): Acute Oral Neurotoxicity Screening Study in Rats: Lab Project Numbers: 23338: T3050251: 106387. Study prepared by Bayer AG, Unpublished.
- FAO/WHO 2002 Manual on Development and Use of FAO and WHO Specifications for Pesticides, 1st edition, FAO, Rome, 2002.
- Flucke 1980 Flucke W., (1980) Boe 5812315 (propoxur). Acute toxicity studies. Report No. 9295 dated July 8, 1980 from Bayer AG Institut für Toxikologie, Wuppertal-Elberfeld, Unpublished.
- Grohs 2004a Bayer_Minutes_Closed_Meeting_JMPS_2003_Answer BCS.doc, attachment to e-mail from R. Grohs (BCS) to G. Vaagt (FAO) and M. Zaim, dated 2 February 2004.
- Grohs 2004b E-mail from R. Grohs (BCS) to G. Vaagt (FAO) and M. Zaim, dated 17 October 2004.
- Hahnemann & Rühl-Fehlert 1988a Hahnemann S. & Rühl-Fehlert C., (1988a) BOQ 5812315 (common name: propoxur). Chronic feeding test on Syrian gold hamsters (species sensitivity). Report No. 16801 dated June 15, 1988 from Bayer AG Toxicology Division Wuppertal, Unpublished.
- Hahnemann & Rühl-Fehlert 1988c Hahnemann S. & Rühl-Fehlert C., (1988c) BOQ 5812315 (common name: propoxur). Chronic feeding test on NMRI mice (species sensitivity). Report No. 16803 dated June 15, 1988 from Bayer AG Toxicology Division Wuppertal, Unpublished.
- Hahnemann & Rühl-Fehlert 1988d Hahnemann S. & Rühl-Fehlert C., (1988d) BOQ 5812315 (common name: propoxur). Sub-chronic feeding test on female Wistar rats (effect of feed quality). Report No. 16897 dated July 13, 1988 from Bayer AG Toxicology Division Wuppertal, Unpublished.

Hahnemann & Rühl-Fehlert 1988f	Hahnemann S. & Rühl-Fehlert C., (1988f) BOQ 5812315 (common name: propoxur). Chronic feeding test on female Wistar rats over 2 years (dose-effect-time relationship). Report No. 16980 dated August 2, 1988 from Bayer AG Toxicology Division Wuppertal, Unpublished.
Heimann, 1982a	Heimann, K.G. (1982a) Propoxur (the active ingredient of ^R Baygon and Uden ^R). Study of sensitization effect on guinea pigs. Report No. 11218 dated October 15, 1982 from Bayer AG Institut für Toxikologie, Wuppertal-Elberfeld, Unpublished.
Heimann, 1983	Heimann, K.G. (1983) Carbamate UN, technical (c.n. propoxur, Uden active ingredient; Baygon active ingredient) Subacute study on rats compared with carbamate UN, recrystallised.. Report No. 11621 dated March 8, 1983 from Bayer AG Institut für Toxikologie, Wuppertal-Elberfeld, Unpublished.
Herbold, 1980a	Herbold, B. (1980a) BOE 5812315, dominant lethal study on male study on male mouse to test for mutagenic effects. Report No. 8808 dated January 7, 1980 from Bayer AG, Institute of Toxicology, Wuppertal-Elberfeld, Unpublished.
Herbold, 1980b	Herbold, B. (1980b) BOE 5812315, micronucleolus test on mouse to evaluate BOE 5812315 for mutagenic potential. Report No. 9274 dated June 27, 1980 from Bayer AG, Institute of Toxicology, Wuppertal-Elberfeld, Unpublished.
Hobik, 1967	Hobik, H.P. (1967) Histological investigations on the spinal cord and nervi ischiadici from neurotoxicity tests on hens with Bayer 39007 Report No. 373 of July 10, 1967, Institute of Experimental Pathology, Bayer AG.
Hoffmann & Gröning, 1984	Hoffmann, K. & Gröning P. (1984) BOQ 5812315 (BOE 5812315, c.n. propoxur) Chronic toxicity to dogs on oral administration (12-months feeding study). Report No. 12605 dated April 11, 1984 from Bayer AG Institute of Toxicology, Wuppertal-Elberfeld, Unpublished.
Hoffmann & Rühl, 1985	Hoffmann, K. & Rühl, Chr. (1985) Propoxur (BOQ 5812315). Subchronic study of toxicity to Rhesus monkeys after oral administration by stomach tube for 13 weeks to check for possible findings in the urinary bladder. Report No. 13799 dated August 27, 1985 from Bayer AG Institute of Toxicology, Wuppertal-Elberfeld, Unpublished.
Inukai & Iyatomi 1978	Inukai, H. & Iyatomi, A. (1978) Propoxur. Mutagenicity test on bacterial systems. Report No. 103 dated February 24, 1978 from Nitokuno Agricultural Chemicals Institute, Toyoda, Japan, Unpublished.
JMPR 1989	Evaluations of pesticide residues in food -1989 (Part II, Toxicology). FAO Plant Production and Protection Paper 100/2, FAO, Rome, 1990
Kimmerle & Iyatomi, 1976	Kimmerle, G. & Iyatomi, A., (1976) Toxicity of propoxur to rats by subacute inhalation. Jap. J. Ind. Health, 18: 375-382.
Kimmerle & Solmecke 1971	Kimmerle, G. & Solmecke, B. (1971) Subacute dermal application to rabbits, report No. 2528 of January 19, 1971, Institute of Toxicology, Bayer AG.
Kimmerle, 1966a	Kimmerle, G. (1966a) Neurotoxic studies on active ingredient BOE 5812315, Report of March 03, 1966, Institute of Toxicology, Bayer AG.
Lamb 1981	Lamb, D. (1981) Acute Dietary LC ₅₀ of Technical Propoxur (Baygon) to Bobwhite Quail: Study No. 80-175-06: Mobay Report No. 80046. Study prepared by Mobay Chemical Corp., Unpublished.
Patterson 1980	Patterson, D.R. (1980) Histopathology Report on BOE 5812315 Mouse Study, Report No. 2317-262/13 of May 1980, Hazleton Laboratories, Harrogate.
Pauluhn 1988	Pauluhn, J. (1988) BOQ 5812315 (common name: propoxur) study of the inhalation toxicity in accordance with OECD Guideline No. 403. Report No. 16966 dated July, 1988 from Bayer AG Institute of Toxicology, Wuppertal, Unpublished.
Pauluhn 1993	Pauluhn, J. (1993) Untersuchungen zur akuten Inhalationstoxizität an der Ratte nach OECD Guideline Nr. 403, Report No. 22224 of April 30, 1993, Fachbereich Toxikologie, Bayer AG.
Riegner 2005	Riegner K., 2005-03-02. Propoxur TGA1: A comparison of the purities and impurity profiles of 'Carbamate UN umkr. techn.' and "Propoxur KCC techn." Bayer CropScience, Monheim. Unpublished.

- Sethi 2005 Sethi S.P., undated. Comparison of manufacturing process information, specifications and batch analyses for Propoxur active constituent submitted to WHO with the information submitted to APVMA. Report attached to e-mail sent by Sethi S.P. to Zaim M., WHO, on 26 August, 2005.
- Sheets & Fuss 1991 Sheets, L.; Fuss, M. (1991) Primary Dermal Irritation Study with Technical Grade Propoxur (Baygon) in Rabbits: Lab Project Number 90-325-HA. Study prepared by Mobay Corp, Unpublished.
- Shirasu *et al.*, 1976 Shirasu, Y., Moriya, M., Kato, K., Furuhashi, A. & Kada, T. (1976) Mutagenicity screening of pesticides in the microbial system. *Mutat. Res.* 40: 19-30.
- Siebert & Eisenbrand, 1974 Siebert, D. & Eisenbrand, E. (1974) Induction of mitotic gene conversion in *Saccharomyces cerevisiae*. *Mutat Res.* 22: 121-126.
- Siebert & Lemperle, 1974 Siebert, D. & Lemperle, E. (1974) Genetic effects of herbicides: induction of mitotic gene conversion in *Saccharomyces cerevisiae*. *Mutat. Res.* 22: 111-120.
- Sturdivant & Halliburton 1998 Acute oral toxicity study with Baygon technical in rats, Bayer Corporation, Stilwell, KS, USA. Unpublished.
- Suter 1990 Suter, P., Biedermann, K. & Luetkemeier, H. (1990) BOQ 5812315 (c.n. Propoxur): Two-Generation Reproduction Study in the Rats: Report 100650. Prepared by Research and Consulting Co., Ag.; Umweltchemie Ag, Unpublished.
- Thyssen 1978 Thyssen, J. (1978) Propoxur-Untersuchungen an der Haut und am Auge von Kaninchen. Report dated September 19, 1978 from Bayer AG Institute of Toxicology, Wuppertal-Elberfeld, Unpublished.
- USEPA 1997 Reregistration Eligibility Decision on Propoxur, EPA, Washington DC, USA, 1997.
- WHO 2000 The WHO Recommended Classification of Pesticides by Hazard and Guidelines to Classification 2000-2002, Geneva. (Document WHO/PCS/01.5).
- Yamane 1986b Yamane, S. (1986) Primary eye irritation study of propoxur in rabbits. Report No. D-0736 (DT-10) dated September 12, 1986 from Hita Research Laboratories, Japan.